

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
29 March 2001 (29.03.2001)

PCT

(10) International Publication Number
WO 01/21807 A1(51) International Patent Classification⁷: C12N 15/40,
5/10, 7/01, C07K 14/18, 16/10, A61K 31/70, 39/29H. [US/US]; 17517 White Ground Road, Boyds, MD
20841 (US).

(21) International Application Number: PCT/US00/25987

(74) Agent: FEILER, William, S.; Morgan & Finnegan,
L.L.P., 345 Park Avenue, New York, NY 10154 (US).

(22) International Filing Date:

22 September 2000 (22.09.2000)

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

60/155,823 23 September 1999 (23.09.1999) US

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).(71) Applicant (*for all designated States except US*): THE
GOVERNMENT OF THE UNITED STATES OF
AMERICA as represented by THE SECRETARY,
DEPARTMENT OF HEALTH SERVICES [US/US];
National Institutes of Health, Office of Technology Trans-
fer, Suite 325, 6011 Executive Boulevard, Rockville, MD
20852 (US).**Published:**

- With international search report.
- Before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments.

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): FORNS, Xavier
[US/US]; Apartment 1106, 4400 East-West Highway,
Bethesda, MD 20814 (US). BUKH, Jens [DK/US];
2018 Baltimore Road, #J42, Rockville, MD 20851 (US).
EMERSON, Suzanne, U. [US/US]; 9903 Parkwood
Drive, Bethesda, MD 20814 (US). PURCELL, Robert,For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

WO 01/21807 A1

(54) Title: HEPATITIS C VIRUS ENVELOPE TWO PROTEIN (E2) WHICH LACKS ALL OR PART OF THE HYPERVARIABLE REGION ONE (HVR1), CORRESPONDING NUCLEIC ACIDS, CHIMERIC VIRUSES AND USES THEREOF

(57) Abstract: The present invention relates to nucleic acid molecules encoding hepatitis C virus, chimeric hepatitis C virus or hepatitis C virus envelope two protein which lacks all or part of hypervariable region one of the envelope two protein. The invention further relates to the use of these nucleic acid molecules and their encoded polypeptides as vaccine candidates.

Title of Invention

**HEPATITE C VIRUS ENVELOPE TWO PROTEIN (E2) WHICH LACKS ALL
OR PART OF THE HYPERVARIABLE REGION ONE (HVR1) ,
CORRESPONDING NUCLEIC ACIDS, CHIMERIC VIRUSES AND USES
THEREOF**

Field of Invention

The present invention relates to nucleic acid molecules that encode a hepatitis C virus (HCV) envelope two protein which lacks all or part of the hypervariable region one (HVR1) of the envelope two (E2) protein. The invention further relates to the use of the nucleic acid molecules and their encoded polypeptides as vaccine candidates.

Background of Invention

Hepatitis C virus (HCV) is a positive-sense single-strand RNA virus belonging to the *Flaviviridae* family of viruses (Rice, 1996).

At present, more than 80% of individuals infected with HCV become chronically infected and these chronically infected individuals have a relatively high risk of developing chronic hepatitis, liver cirrhosis and hepatocellular carcinoma (Hoofnagle, 1997). The only effective therapy for chronic hepatitis C, interferon (IFN), alone or in combination with ribavirin, induces a sustained response in less than 50% of treated patients (Davis et al., 1998; McHutchinson et al., 1998). Consequently, HCV is currently the most common cause of end-stage liver failure and the reason for about 30% of liver transplants performed in the U.S. (Hoofnagle, 1997). As a result of the inability to develop a universally effective therapy against HCV infection, it is estimated that there are still more than 25,000 new infections yearly in the U.S. (Alter

1997) Moreover, since there is no vaccine for HCV, HCV remains a serious public health problem.

A remarkable characteristic of HCV is its genetic heterogeneity, which is manifested throughout the genome (Bukh et al., 1995). HCV circulates as a quasispecies of closely related genomes in an infected individual. Globally, six major HCV genotypes (genotypes 1-6) and multiple subtypes (a, b, c, etc.) have been identified (Bukh et al., 1993; Simmonds et al., 1993). The nucleotide and deduced amino acid sequences among isolates within a quasispecies generally differ by < 2%, whereas those between isolates of different genotypes vary by as much as 35%.

The most heterogeneous regions of the genome are found in the two envelope genes E1 and E2; in particular, the hypervariable region 1 (HVR1) at the N-terminus of E2 (Hijikata et al., 1991; Weiner et al., 1991).

The fact that the HVR1 is the region of the genome with the highest degree of genetic variability suggests that it is under strong immune pressure. Indeed, Ray et al. (1999) recently reported that patients who developed a chronic infection had a higher rate of non-synonymous mutations within the HVR1 as compared with the E1 protein while the reverse was observed in patients who were able to clear the infection. These authors therefore hypothesized that the HVR1 region of HCV might act as a decoy antigen by stimulating a strong immune response that is ineffective in clearing viremia.

However, the proposed role of the HVR1 as an immunologic decoy is not easily reconciled with prior studies in which a hyperimmune rabbit serum raised against the HVR1 was demonstrated to be capable of neutralizing HCV *in vitro* (Farci et al., 1996, Shimizu et al., 1996). Moreover, the presence of amino acids in the carboxy-terminal half of HVR1 which are

conserved across genotypes suggests that HVR1 might be required for HCV replication.

Summary of Invention

The present invention relates to nucleic acid molecules which lack all or part of the coding sequence of the HVR1 region of the envelope 2 (E2) gene of hepatitis C virus (HCV).

In the first embodiment, the nucleic acid molecule of the invention comprises the genome of an infectious hepatitis C virus in which the HVR1 of the envelope 2 gene of the infectious HCV has been deleted.

It is therefore an object of the invention to provide nucleic acid sequence which encodes infectious hepatitis C virus lacking HVR1. Such nucleic acid sequence is referred to throughout the application as "ΔHVR1-infectious nucleic acid sequence."

In a second embodiment, the nucleic acid molecule comprises chimeric genomes of chimeric hepatitis C viruses in which the structural region (core and envelope genes) or the envelope gene of a pestivirus genome, [for example, a bovine viral diarrhea virus (BVDV)] or a flavivirus genome (for example, a dengue virus or a yellow fever virus) are replaced by the corresponding structural region or E1 and E2 genes of an HCV in which the HVR1 region of HCV E2 has been removed. It is therefore an object of the invention to provide nucleic acid sequence which encodes chimeric HCV lacking the HVR1 of the HCV E2 gene. Such nucleic acid sequence is referred to throughout the application as "ΔHVR1-chimeric nucleic acid sequences."

The present invention also relates to the in vitro and in vivo production of ΔHVR1-infectious HCV or ΔHVR1-chimeric HCV

viruses from the Δ HVR1-infectious or Δ HVR1-chimeric nucleic acid sequences of the invention.

The present invention also relates to the use of the Δ HVR1 viruses of the invention to identify cell lines capable of supporting the replication of the viruses.

The invention also relates to the use of the Δ HVR1 infectious or chimeric nucleic acid sequences of the invention in the production of Δ HVR1-infectious HCV or Δ HVR1-chimeric HCV respectively, and the use of these virions for the development of inactivated or attenuated vaccines to prevent HCV in a mammal.

In a third embodiment, the DNA construct comprises an HCV E2 gene lacking the HVR1 region. Such a " Δ HVR1-E2 gene" may also be linked in tandem in the DNA construct with an HCV E1 gene. When contained alone or in tandem with the E1 gene, the Δ HVR1-E2 gene may be further modified at its carboxy-terminus to produce either a secreted or surface expressed Δ HVR1-E2 protein.

The invention further relates to pharmaceutical compositions and DNA-based vaccines which comprise the nucleic acid molecules of the invention.

The invention also relates to methods of preventing or treating HCV in a mammal comprising administering the nucleic acid molecules of the invention to a mammal in an amount effective to stimulate the production of a protective humoral and/or cellular immune response to HCV.

The invention also provides a kit for the treatment or prevention of HCV, the kit comprising a DNA molecule of the invention useful as an immunogen in generating a protective immune response to HCV.

The invention further relates to the use of the nucleic acid molecules of the invention as immunogens to

generate antibodies to the Δ HVR1-infectious HCV, the Δ HVR1-chimeric HCV or the Δ HVR1-E2 protein, preferably neutralizing antibodies. The invention therefore relates to the use of such antibodies in passive immunoprophylaxis and to pharmaceutical compositions which comprise these antibodies.

The invention also relates to transformation of host cells with nucleic acid molecules of the invention to produce host cells which express Δ HVR1-infectious HCV, the Δ HVR1-chimeric HCV or the Δ HVR1-E2 protein.

The invention further relates to the use of host cells expressing Δ HVR1-infectious HCV, the Δ HVR1-chimeric HCV or the Δ HVR1-E2 protein as immunogens to stimulate a protective immune response to HCV.

The present invention, of course, also relates to Δ HVR1 E2 protein produced from the Δ HVR1-E2 gene constructs of the invention or obtained from the Δ HVR1-infectious HCVs or Δ HVR1-chimeric HCVs of the invention. These Δ HVR1-E2 proteins may be used as vaccines for immunizing mammals, especially humans, against HCV.

Brief Description Of Figures

Figures 1A-1F show the nucleotide sequence (SEQ ID NO: 1) of the infectious hepatitis C virus clone of genotype 1a [H77C(Δ HVR1)] which lacks the hypervariable region one (HVR1) of the second envelope protein and Figures 1G-1H show the amino acid sequence (SEQ ID NO: 2) encoded by the clone. The complete sequence of H77C(Δ HVR1) is identical to pCV-H77C but lacks the fragment from nucleotide positions 1491 to 1571 which encodes HVR1 (Yanagi et al., 1997; ATCC accession number PTA-157).

Figure 2 shows *in vitro* transcription-translation of constructs ElE2-715 (Lanes 1, 3, 4, 5, and 6) and Δ HVR1-7 (Lane 8) with rabbit reticulocyte lysates with and without the addition of canine microsomal membranes.

Figure 3 shows the results of qualitative reverse transcriptase-nested polymerase chain reaction (RT-PCR) for HCV-RNA, \log_{10} HCV GE (genome equivalent) titer (in-house RT-PCR and Amplicor HCV Monitor, Roche Diagnostics), second generation ELISA for anti-HCV and serum levels of alanine aminotransferase (ALT) in chimpanzee 1590 following transfection with RNA transcripts of the HCV deletion mutant H77C(Δ HVR1).

Figure 4 shows infection of chimpanzee 96A008 with HCV lacking HVR1. Serum samples were collected weekly from the chimpanzee and monitored for HCV-RNA [in-house RT-nested PCR and HCV Monitor test version 2.0 (Roche)], HCV antibodies (second generation ELISA, Abbott Laboratories) and liver enzyme levels (ALT, Anilytics). PBMC were collected weekly and tested for HCV-specific proliferative capacity (peripheral CD4) with a panel of recombinant HCV proteins [C22 (core), C33-c (NS3), C100 (NS3-NS4), and NS5]. The peripheral CD8+ T cell response (peripheral CTL) was tested by stimulating PBMC with a large panel of HCV peptides corresponding to known CTL epitopes. Expanded T cells isolated from liver biopsy samples were tested for HCV-specific proliferative responses (Intrahepatic CD4) +, positive; - negative. Liver biopsies were examined also for necroinflammatory changes [0 (normal), 1+, 2+, 3+, 4+]. At week 0, the chimpanzee was inoculated intravenously with 90 ml of plasma from chimpanzee 1590 (week 4 after transfection). At a titer of 10 GE/ml this represented an inoculum of approximately 900 genome equivalents of HCV.

Detailed Description of The Invention

The present inventors surprisingly observed that HCV cDNA from which the HVR1 of the E2 gene had been deleted, encoded a hepatitis C virus which was able to replicate in vivo and stimulated a strong cellular immune response. Figure 1A-1F shows the nucleic acid sequence of this infectious HCV clone which was constructed using pCV-H77C (ATCC accession number PTA-157), an infectious HCV clone of genotype 1a.

The present invention therefore relates in one embodiment to a nucleic acid molecule which comprises the genome of an infectious hepatitis C virus in which the HVR1 of the envelope 2 gene of the infectious HCV has been deleted.

The E2 protein consists of amino acid 384 to 746 of the HCV polyprotein and the HVR1 of the E2 protein consists of amino acid 384 to 410.

It is therefore an object of the invention to provide nucleic acid sequence which encodes infectious hepatitis C virus lacking HVR1. Such nucleic acid sequence is referred to throughout the application as "ΔHVR1-infectious nucleic acid sequence."

It is understood that the deletion of HVR1 of the E2 gene in the HCV genome can be made in infectious HCV clones of any genotype. For example, infectious HCV clones of different genotypes which have been constructed include those of Kolykhalov et al., (1997) and Yanagi et al. (1997, 1998) [who reported the derivation from HCV strains H77 (genotype 1a) and HC-J4 (genotype 1b) of cDNA clones of HCV that are infectious for chimpanzees], and Yanagi et al. (Yanagi, 1999) [who reported the construction of an infectious cDNA clone from HCV strain HC-J6 (genotype 2a)]. Preferably, the infectious HCV clones are infectious HCV clones of genotype 1a (ATCC accession number PTA-

157), 1b (ATCC accession number 209596) or 2a (ATCC accession number PTA-153).

In a second embodiment, the nucleic acid molecule of the invention comprises the genome of a flavivirus or pestivirus in which one envelope gene of the flavivirus or pestivirus is replaced by the E2 gene of an infectious HCV from which the HVR1 region of HCV E2 has been removed.

It is therefore an object of the invention to provide nucleic acid sequence which encodes chimeric HCV lacking the HVR1 of the HCV E2 gene. Such nucleic acid sequence is referred to throughout the application as "ΔHVR1-chimeric nucleic acid sequence."

The *Flaviviridae* family of viruses which may be used to make the ΔHVR1-chimeric nucleic acid sequences include, but are not limited to, dengue virus, bovine viral diarrhea virus, yellow fever virus and Kunjin virus..

Preferably, the ΔHVR1-chimeric nucleic acid sequences of the invention are made using the structural region or E1 and E2 genes of an infectious HCV clone of any genotype. Preferably, the infectious HCV clones are infectious HCV clones of genotype 1a (ATCC accession number PTA-157), 1b (ATCC accession number 209596) or 2a (ATCC accession number PTA-153).

The deletion of HVR1 which may be made in the sequence of the invention is at least 5 amino acids in length, preferably 10 amino acids, and most preferably the entire 27 amino acids of HVR1. The nucleic acid molecules of the invention therefore comprise genomes of HCV which lack at least a fragment of the HVR1 sequence.

Of course, it is understood that in deleting all or part of the HVR1 sequence, one may extend the deletions further in the carboxy-terminal direction of the E2 protein as long as

the resultant nucleic acid molecule is capable of replicating in a chimpanzee and producing the E2 protein lacking all or part of HVR1.

The Δ HVR1-infectious nucleic acid sequence of the invention may further include one or more mutations such as those described in Example 4 which result in amino acid changes. One mutation is located within the E2 gene and results in a change from leucine to histidine at amino acid position 615. Another mutation is located within the NS3 serine-protease domain and results in a change from arginine to histidine at amino acid position 1143. A third mutation is located within the NS5B RNA-polymerase domain and results in a change from glutamic acid to aspartic acid at amino acid position 2875.

The present invention further relates to the in vitro and in vivo production of hepatitis C viruses from the Δ HVR1-infectious nucleic acid sequences of the invention, and the production of chimeric viruses from the Δ HVR1-chimeric nucleic acid sequences of the invention.

In one embodiment, the Δ HVR1-infectious nucleic acid sequences or the Δ HVR1-chimeric nucleic acid sequences of the invention can be inserted into an expression vector that functions in eukaryotic cells. Eukaryotic expression vectors suitable for producing high efficiency gene transfer in vivo are well known to those of ordinary skill in the art and include, but are not limited to, plasmids, vaccinia viruses, retroviruses, adenoviruses and adeno-associated viruses.

The sequences contained in the recombinant expression vector can then be transcribed in vitro by methods known to those of ordinary skill in the art in order to produce RNA transcripts which encode the HCV or chimeric viruses of the invention. The HCV or chimeric viruses of the invention may

then be produced by transfecting cells by methods known to those of ordinary skill in the art with either the in vitro transcription mixture containing the RNA transcripts or with the recombinant expression vectors containing the nucleic acid sequences described herein.

Where transfection of cells with recombinant expression vectors containing the nucleic acid sequences of the invention is used, transfection may be done by methods known in the art such as electroporation, precipitation with DEAE-Dextran or calcium phosphate, or incorporation into liposomes.

Suitable cells or cell lines for culturing the HCV or chimeric viruses of the invention include, but are not limited to, EBTr and Huh7.

The present invention also relates to the use of the Δ HVR1-infectious nucleic acid sequences or the Δ HVR1-chimeric nucleic acid sequences of the invention to identify cell lines capable of supporting the replication of HCV and the chimeric viruses of the invention.

The invention further relates to the use of the Δ HVR1-infectious nucleic acid sequences or the Δ HVR1-chimeric nucleic acid sequences of the invention to develop inactivated or attenuated vaccines to prevent hepatitis C in a mammal. For example, virions from cell lines infected with the HCV or chimeric viruses of the invention, or transfected with a Δ HVR1-infectious nucleic acid sequence or a Δ HVR1-chimeric nucleic acid sequence of the invention, can be purified from the cells and inactivated by methods known to those of ordinary skill in the art. The inactivated virions can be used to immunize mice, and if neutralizing antibody to HCV is produced, the virions can then be used to immunize chimpanzees to determine whether the antibodies are protective. Alternatively, cells infected with

the viruses of the invention may be passaged in cell culture to produce attenuated viruses which can be tested as candidate live vaccines. In assaying the ability of the viruses of the invention to infect mammals one can assay sera or liver of the infected mammal by RT-PCR to determine viral titer. In addition, the virulence phenotype of the virus produced by transfection of mammals with the sequences of the invention can be monitored by methods known in the art such as measurement of liver enzyme levels (alanine aminotransferase (ALT) or isocitrate dehydrogenase (ICD)) or by histopathology of liver biopsies.

When used as a vaccine, the HCV or chimeric virions can be administered alone or in a suitable diluent, including, but not limited to, water, saline, or some type of buffered medium. The vaccine according to the present invention may be administered to an animal, especially a mammal, and most especially a human, by a variety of routes, including, but not limited to, intradermally, intramuscularly, subcutaneously, or in any combination thereof. Of course, it is understood that formulations or compositions comprising the HCV or chimeric virions of the invention may be used either therapeutically or prophylactically to treat or prevent the signs and symptoms of Hepatitis C.

In a third embodiment, the nucleic acid molecule of the invention comprises an HCV E2 gene lacking the HVR1 region. Such a " Δ HVR1-E2 gene" may also be linked in tandem in the DNA construct with an HCV E1 gene.

When contained alone or in tandem with the E1 gene, the Δ HVR1-E2 gene may be further modified at its carboxy-terminus to produce either a secreted or surface expressed Δ HVR1-E2 protein.

A Δ HVR1-E2 gene encoding E2 protein targeted to the cell surface is preferred. Such a construct may be constructed by fusing an endoplasmic reticulum signal sequence to the amino-terminus of the nucleic acid sequence which encodes the truncated E2 gene fused at its carboxy-terminus to a plasma membrane anchor sequence.

By endoplasmic reticulum (ER) signal sequence is meant a nucleic acid sequence which encodes a continuous stretch of amino acids, typically about 15 to about 25 residues in length, which are known in the art to be generally located at the amino terminus of proteins and are capable of targeting proteins to the endoplasmic reticulum. Such ER signal sequences are known to those of skill in the art (see, for example, van Heijne, G. J. Mol. Biol., (1985) 184:99-105) and those of skill in the art would understand that even though their amino acid sequences may vary, such ER signal sequences are functionally interchangeable. Examples of ER signal sequences which may be used in the chimeric genes of the invention include, but are not limited to, the 20-carboxy-terminal amino acids of the full-length HCV E1 protein (amino acids 364-383 of the HCV polyprotein), which serves as the natural signal sequence of the E2 protein or the murine Ig kappa-chain V-J2-C signal peptide sequence contained in the pDisplay vector.

Where the truncated HCV envelope protein is a truncated E2 protein, the approximately 30 carboxy-terminal amino acids of E2 have been identified to contain an ER retention sequence and its removal and replacement with a plasma membrane anchor sequence is believed to be critical for expression of the truncated E2 protein on the cell surface. Thus, the truncated E2 protein contains a truncation of at least the 20 carboxy-terminal amino acids of the full-length E2

protein, more preferably, a truncation of at least the 25 carboxy-terminal amino acids, and most preferably, a truncation of at least about the 30 carboxy-terminal amino acids. By "plasma membrane anchor sequence" as used in the chimeric gene of the invention is meant a nucleic acid sequence which encodes an amino acid sequence that allows for retention of at least part of the protein in the plasma membrane of a cell. At a minimum, a plasma membrane anchor sequence encodes a sequence of hydrophobic amino acids of sufficient length to span the lipid bilayer of the plasma membrane. Such hydrophobic sequences are known in the art as transmembrane domains and are typically found at the carboxy-terminus of many proteins found on the surface of cells or virions. These transmembrane domains are typically at least 20 to 30 amino acids in length and are followed by charged cytoplasmic domains of varying lengths.

It is therefore understood that the plasma membrane anchor sequence encoded by the coding sequence of the invention may contain in addition to a transmembrane domain of a virion or a protein found on the surface of a cell, a cytoplasmic domain.

Perferably, the encoded plasma membrane anchor sequence is at least twenty amino acids in length, more preferably, from about 20 to about 100 amino acids in length, and most preferably, from about 30 to about 70 amino acids in length. Examples of plasma membrane anchor sequences include, but are not limited to, hydrophobic transmembrane domains of receptors such as those for insulin and for a number of growth factors including platelet-derived growth factor (PDGF) and epidermal growth factor (EGF), as well as the transmembrane domains of viral proteins that are anchored in the lipid envelope of the intact virion such as the transmembrane domains of the vesicular stomatitis and rabies virus G proteins.

Preferred plasma membrane anchor sequences for inclusion in the chimeric genes of the invention are sequences which encode the 50 amino acid transmembrane domain of the PDGF receptor as contained in the pDisplay vector described in the Examples, the carboxy-terminal 64 and 37 amino acids respectively of the CD4 and decay accelerating factor (DAF) proteins (these sequences constitute the transmembrane and cytoplasmic domains of the CD4 and DAF proteins) and the 49 carboxy-terminal amino acids of the VSV G protein (also constituting the transmembrane and cytoplasmic domains of the VSV G protein).

Of course, one of ordinary skill in the art would readily understand that other transmembrane domains suitable for use as plasma membrane anchor sequences in the chimeric genes of the invention are known or could be readily identified by carrying out carboxy-terminal deletions of known plasma membrane or viral envelope proteins (see, for example, Men et al (J. Virol. (1991) 65; 1400-1407)).

A Δ HVR1-E2 gene encoding a secreted E2 protein may be constructed by fusing an endoplasmic reticulum signal sequence to the nucleic acid sequence which encodes the truncated E2 gene. In this case, the truncated E2 gene contains a truncation of at least the 31 carboxy-terminal amino acids (amino acids 716-746), and more preferably, a truncation of at least the 85 carboxy-terminal amino acids (amino acids 662-746).

Like the Δ HVR1-infectious nucleic acid sequences and the Δ HVR1-chimeric nucleic acid sequences described above, it is understood that the Δ HVR1-E2 nucleic acid sequences of the invention may lack all or part of the HVR1 sequence. The deletion of HVR1 is at least 5 amino acids in length, preferably 10 amino acids, and most preferably the entire 27 amino acids of

HVR1. The Δ HVR1-E2 nucleic acid sequences of the invention may also contain further truncations at the carboxy-terminus of the E2 gene of HCV, and/or mutations at amino acid position 615 as described above.

The present invention therefore relates to insertion of the nucleic acid molecules comprising the Δ HVR1-E2 gene of the invention into a suitable expression vector that functions in eukaryotic cells, preferably in mammalian cells. By suitable it is meant that the vector is capable of carrying and expressing a chimeric gene of the invention. The expression vector therefore contains at least one promoter and any other sequences necessary or preferred for appropriate transcription and translation of the Δ HVR1-E2 gene. Preferred expression vectors include, but are not limited to, plasmid vectors.

The invention also relates to the use of expression vectors containing the Δ HVR1-E2 nucleic acid molecules of the invention as immunogens to produce protective antibodies to HCV. Direct transfer of the Δ HVR1-E2 nucleic acid sequences of the invention to a mammal, preferably a primate, more preferably a human, may be accomplished by injection by needle or by use of other DNA delivery devices such as the gene gun. Possible routes of administration of the expression vector include, but are not limited to, intravenous, intramuscular, intradermal, subcutaneous, intraperitoneal and intranasal.

Since the existence of different genotypes with a low degree of homology within the envelope proteins diminishes the hope of identifying conserved neutralization epitopes (Bukh J, et al, Sem Liver Dis (1995); 15:41-63), it is likely that a polyvalent vaccine will be needed to generate broadly reactive neutralizing antibodies. Thus, in a preferred embodiment, nucleic acid molecules comprising Δ HVR1-E2 nucleic acid

sequences of isolates from multiple genotypes of HCV may be administered together to provide protection against challenge with multiple genotypes of HCV.

Accordingly, those of ordinary skill in the art would readily understand that multiple copies of different Δ HVR1-E2 nucleic acid sequences may be inserted into a single vector such that a host cell transformed or transfected with the vector will produce multiple envelope proteins. For example, a polycistronic vector in which multiple different Δ HVR1-E2 genes may be expressed from a single vector is created by placing expression of each gene under control of an internal ribosomal entry site (IRES) (Molla, a. et al. Nature, 356:255-257 (1992); Gong, S.K. et al. J. of Virol., 263:1651-1660 (1989)). In a preferred embodiment, copies of different Δ HVR1-E2 nucleic acid sequences are inserted into multiple vectors and transformed or transfected into host cells so that multiple envelope proteins can be produced.

The expression vectors containing the Δ HVR1-E2 nucleic acid sequences of the invention may be supplied in the form of a kit, alone, or in the form of a pharmaceutical composition.

Suitable amounts of material to administer for prophylactic and therapeutic purposes will vary depending on the route selected and the immunogen (E2 gene with different deletions) administered. One skilled in the art will appreciate that the amounts to be administered for any particular treatment protocol can be readily determined without undue experimentation. The vaccines of the present invention may be administered once or periodically until a suitable titer of anti-HCV antibodies appear in the blood. A suitable amount of expression vector to be used for prophylactic purposes might be expected to fall in the range of from about 1 μ g to about 5 mg,

more preferably from about 100 µg to about 5 mg, and most preferably from about 1 mg to about 2 mg. Such administration will, of course, occur prior to any sign of HCV infection. Further, one of skill in the art will readily understand that the amount of vector to be used will depend on the size and species of animal the vector is to be administered to.

A vaccine of the present invention may be employed in sterile liquid forms such as solutions or suspensions. Any inert carrier is preferably used, such as saline or phosphate-buffered saline, or any such carrier in which the expression vector of the present invention can be suitably suspended. The vaccines may be in the form of single dose preparations or in multi-dose flasks which can be utilized for mass-vaccination programs of both animals and humans. Of course, specific adjuvants such as CpG motifs (Krieg, A.K. et al. (1995) Nature 374:546 and Krieg et al. (1996)) J. Lab. Clin. Med., 128:128) may prove useful with DNA-based vaccines or other vaccines.

The DNA-based vaccines will normally exist as physically discrete units suitable as a unitary dosage for animals, especially mammals, and most especially humans, wherein each unit will contain a predetermined quantity of active material calculated to produce the desired immunogenic effect in association with the required diluent. The dose of said vaccine or inoculum according to the present invention is administered at least once. In order to increase the antibody level, a second or booster dose may be administered at some time after the initial dose. The need for, and timing of, such booster dose will, of course, be determined within the sound judgment of the administrator of such vaccine or inoculum and according to sound principles well known in the art. For example, such booster dose could reasonably be expected to be advantageous at

some time between about 2 weeks to about 6 months following the initial vaccination. Subsequent doses may be administered as indicated.

The Δ HVR1-E2 nucleic acid sequences of the present invention can also be administered for purposes of therapy, where a mammal, especially a primate, and most especially a human, is already infected, as shown by well-known diagnostic measures. When expression vectors containing the chimeric genes of the present invention are used for such therapeutic purposes, much of the same criteria will apply as when they are used as a vaccine, except that inoculation will occur post-infection. Thus, when the expression vectors of the present invention are used as therapeutic agents in the treatment of infection, the therapeutic agent comprises a pharmaceutical composition containing a sufficient amount of the expression vector so as to elicit a therapeutically effective response in the organism to be treated. Of course, the amount of pharmaceutical composition to be administered will, as for vaccines, vary depending on the immunogen contained therein and on the route of administration.

The therapeutic agent according to the present invention can thus be administered by subcutaneous, intramuscular, intradermal or intranasal routes. One skilled in the art will certainly appreciate that the amounts to be administered for any particular treatment protocol can be readily determined without undue experimentation. Of course, the actual amounts will vary depending on the route of administration as well as the sex, age, and clinical status of the subject which, in the case of human patients, is to be determined with the sound judgment of the clinician.

The therapeutic agent of the present invention can be employed in sterile liquid forms such as solutions or

suspensions. An inert carrier is preferably used, such as saline, phosphate-buffered saline, or any such carrier in which the expression vectors of the present invention can be suitably suspended. The therapeutic agents may be in the form of single dose preparations or in multi-dose flasks, which can be utilized for mass-treatment programs of both animals and humans. Of course, when the expression vectors of the present invention are used as therapeutic agents, they may be administered as a single dose or as a series of doses, depending on the situation as determined by the person conducting the treatment.

The invention also relates to transformation of host cells with nucleic acid molecules of the invention to produce host cells which express HVR1-infectious HCV, the Δ HVR1-chimeric HCV or the Δ HVR1-E2 protein.

The invention further relates to the use of host cells expressing Δ HVR1-infectious HCV, the Δ HVR1-chimeric HCV or the Δ HVR1-E2 protein as immunogens to stimulate a protective immune response to HCV.

The present invention, of course, also relates to Δ HVR1 E2 protein produced from the Δ HVR1-E2 gene constructs of the invention or obtained from the Δ HVR1-infectious HCVs or Δ HVR1-chimeric HCVs of the invention.

The nucleic acids, polypeptides and viruses of the present invention can also be utilized in the production of antibodies against HCV. The term "antibody" is herein used to refer to immunoglobulin molecules and immunologically active portions of immunoglobulin molecules. Examples of antibody molecules are intact immunoglobulin molecules, substantially intact immunoglobulin molecules and portions of an immunoglobulin molecule, including those portions known in the

art as Fab, F(ab')₂ and F(v) as well as chimeric antibody molecules.

Thus, the polypeptides, viruses and nucleic acid sequences of the present invention can be used in the generation of antibodies that immunoreact (i.e., specific binding between an antigenic determinant-containing molecule and a molecule containing an antibody combining site such as a whole antibody molecule or an active portion thereof) with antigenic determinants on the surface of hepatitis C virus particles.

The present invention therefore also relates to antibodies produced following immunization with the nucleic acid sequences, viruses or polypeptides of the present invention. These antibodies are typically produced by immunizing a mammal with an immunogen or vaccine to induce antibody molecules having immunospecificity for polypeptides or viruses produced in response to infection with the nucleic acid sequences of the present invention. When used in generating such antibodies, the nucleic acid sequences, viruses, or polypeptides of the present invention may be linked to some type of carrier molecule. The resulting antibody molecules are then collected from said mammal. Antibodies produced according to the present invention have the unique advantage of being generated in response to authentic, functional polypeptides produced according to the actual cloned HCV genome.

The antibody molecules of the present invention may be polyclonal or monoclonal. Monoclonal antibodies are readily produced by methods well known in the art. Portions of immunoglobulin molecules, such as Fabs, as well as chimeric antibodies, may also be produced by methods well known to those of ordinary skill in the art of generating such antibodies.

The antibodies according to the present invention may also be contained in blood, plasma, serum, hybridoma supernatants, and the like. Alternatively, the antibody of the present invention is isolated to the extent desired by well known techniques such as, for example, using DEAE Sephadex. The antibodies produced according to the present invention may be further purified so as to obtain specific classes or subclasses of antibody such as IgM, IgG, IgA, and the like. Antibodies of the IgG class are preferred for purposes of passive protection.

The antibodies of the present invention are useful in the prevention and treatment of diseases caused by hepatitis C virus in animals, especially mammals, and most especially humans.

In providing the antibodies of the present invention to a recipient mammal, preferably a human, the dosage of administered antibodies will vary depending on such factors as the mammal's age, weight, height, sex, general medical condition, previous medical history, and the like.

In general, it will be advantageous to provide the recipient mammal with a dosage of antibodies in the range of from about 1 mg/kg body weight to about 10 mg/kg body weight of the mammal, although a lower or higher dose may be administered if found desirable. Such antibodies will normally be administered by intravenous or intramuscular route as an inoculum. The antibodies of the present invention are intended to be provided to the recipient subject in an amount sufficient to prevent, lessen or attenuate the severity, extent or duration of any existing infection.

The invention also provides that the nucleic acid molecules, viruses, polypeptides and antibodies of the invention may be supplied in the form of a kit, alone or in the form of a pharmaceutical composition.

All scientific publication and/or patents cited herein are specifically incorporated by reference. The following examples illustrate various aspects of the invention but are in no way intended to limit the scope thereof.

EXAMPLES

Materials and Methods

Construction of An Expression Vector Δ HVR1-7 Containing E1 and a Carboxy-Terminal Truncated E2 Which Lacks the HVR1 Region and the Carboxy-Terminal 31 Amino Acids of E2

Deletion of the nucleotide sequence encoding the HVR1 region was performed by fusion PCR using pCV-H77C, the infectious cDNA clone of HCV genotype 1a (Yanagi et al., 1997; ATCC accession number PTA-157).

Table 1Primers used for the construction of ΔHVR1*

<u>PRIMER</u>	<u>SEQUENCE</u>
SEQ: ID NO:3	
E2-PstI:	5'-ACG CGT <u>CTG CAG</u> CTT AAT GGC CCA GGA CGC GAT GCT TG-3'
SEQ: ID NO:4	
E1-BglII:	5'- ACG CGT <u>AGA TCT</u> TAC CAA GTG CGC AAT TCC TCG GGG-3'
SEQ: ID NO:5	
E1-383R (+10):	5'-TCA GTT GGA TAG CGT CGA CGC CGG CAA ATA GCA G-3'
SEQ: ID NO:6	
E2-411S (+10):	5'-CGT CGA CGC GAT CCA ACT GAT CAA CAC CAA CGG C-3'
SEQ: ID NO:7	
HVR1-mutS:	5'-CTT GTA CCA TCA ATT ACA CCA TAT TC-3'
SEQ: ID NO:8	
HVR1-mutR:	5'-GAT AGT GCC AAT GCC TAT ACG GG-3'
SEQ: ID NO:9	
FUSION 1:	5'-CGT ATA GGC ATT GGC ACT ATC CTT GTA CCA TCA ATT ACA CC-3'
SEQ: ID NO:10	
FUSION2:	5'-GGT GTA ATT GAT GGT ACA AGG ATA GTG CCA ATG CCT ATA CG-3'
SEQ: ID NO:11	
E2-364 NotI:	5'-TTT TTT TTG <u>CGG CCG</u> CAT GGT GGG GAA CTG GGC GAA GGT CC-3'
SEQ: ID NO:12	
E2-661-HindIII:	5'-ACG CGT <u>AAG CTT</u> CTA TTA CTC GGA CCT GTC CCT GTC TTC CAG-3'

*Restriction sites within primers are underlined

In the first step, the sequences encoding the E1 protein (aa 192-383) and a truncated E2 protein (aa 411-715) lacking the HVR1 region (aa 384-410) were PCR amplified from pCV-H77C. Briefly, 50 ng of cDNA were added to a master mix containing 4 μ l of 5X Advantage KlenTaq Buffer, 1.25 μ l of dNTP (10 mM), 1 μ l of 10 μ M sense primer [E1-BglII (SEQ ID NO: 4) for E1 and E2-411S(+10) (SEQ ID NO: 6) for E2], 1 μ l of 10 μ M antisense primer [E1383R (+10) for E1 and E2-PstI for E2] and 1 μ l of the Advantage KlenTaq polymerase mixture. Cycling conditions were 99°C for 1 min followed by 25 cycles of 99°C for 35 sec, 67°C for 30 sec, 68°C for 3 min 30 sec. After purification of the PCR products, a fusion PCR was performed. Briefly, 2.5 μ l of each PCR product were added to a master mix containing 10 μ l of 10X Pfu buffer, 2 μ l of 10 mM dNTP, 5 μ l of 10 μ M sense primer (E1-BglII) (SEQ ID NO: 4), 5 μ l of 10 μ M antisense primer (E2-PstI) (SEQ ID NO: 3), 1 μ l of Pfu and 72 μ l of H₂O. Cycling conditions were 95 °C for 1 min, 67 °C for 1 min, 72 °C for 3 min 30 sec (30 cycles). The fusion PCR product was digested with *Bgl*II and *Pst*I and cloned into the expression vector pDisplay (Invitrogen). A clone with an insert of the correct size (Δ HVR1-7) was selected and sequenced. Sequence analysis of both strands of DNA confirmed that this clone contained the expected sequence of E1 and a truncated E2 lacking the HVR1, i.e., Δ HVR1-7 encodes amino acids 192-383 of the E1 protein fused to amino acids 411-715 of the E2 protein.

Construction of An Infectious clone of HCV Lacking HVR1

Construction of an infectious cDNA clone that was full-length except for the HVR1 was performed by digestion of the expression vector Δ HVR1-7 with *Mun*I (which cuts at

nucleotide positions 1254 and 1983 of pCV-H77C) and cloning of the resulting fragment into the digested HCV cDNA clone H77C. Briefly, a 3 µg sample of the infectious HCV cDNA clone H77C was digested with *MunI* for 3 hours and the enzyme was then inactivated at 65°C for 20 min. The digested cDNA clone, lacking a portion of the E1 and E2 regions (nucleotides 1255 to 1983 of pCV-H77C), was dephosphorylated with calf intestinal alkaline phosphatase. A sample of ΔHVR1-7 was then digested with *MunI* for 3 hours. After inactivation of the *MunI* enzyme, the digested fragment was ligated into the digested pCV-H77C using standard procedures (Forns et al., 1999). A clone containing the correct insert was selected, retransformed and large-scale plasmid DNA was prepared as previously described (Yanagi et al., 1997). The complete sequence of the HVR1 deletion mutant [H77C(ΔHVR1)] was the expected one, that is, identical to pCV-H77C but lacking the fragment from nucleotide positions 1491 to 1571 which encodes HVR1.

In vitro transcription-translation analysis

Non-linearized plasmid ΔHVR1-7 and the control plasmid E1E2-715 were used for *in vitro* transcription-translation. The ΔHVR1-7 plasmid contained the nucleotide sequence encoding E1 (aa 192-383) and E2 (aa 411-715) inserted between a leader sequence which targeted the HCV proteins to the secretory pathway and the transmembrane domain of the platelet-derived growth factor receptor (PDGFR) which anchored the HCV E2 proteins to the plasma membrane in the expression vector pDisplay (Invitrogen). The control plasmid contained the nucleotide sequence encoding E1 (aa 192-383) a carboxy-truncated form of E2 (aa 384-715) of pCV-H77C inserted between a leader sequence which targeted the HCV proteins to the secretory pathway and the transmembrane domain of PDGFR which anchored the

HCV E2 proteins to the plasma membrane in the expression vector pDisplay. Reactions were performed in 25 μ l of the TNT Coupled Reticulocyte Lysate System (Promega) containing [S^{35}] methionine, with or without the addition of canine microsomal membranes at 30 °C for 90 min. Total translation products were separated in 12% SDS/PAGE and identified by autoradiography.

Analysis of cell surface-expression of the E2 protein

Huh7 cells grown in 4-well tissue culture chambers were transfected with plasmids E1E2-715 and Δ HVR1-7 described above. Immuno-fluorescence analysis was performed 48 hours after transfection. Live cells were incubated for 30 minutes with a 1:100 dilution of rabbit hyperimmune serum (*Lmf86*) raised against a peptide encoding the carboxy-terminal 21 amino acids of HVR1 (aa 390-410 of pCV-H77C) or a rabbit hyperimmune serum (*FOR-1*) raised against a peptide within E2, but outside the HVR1 (aa 517-535 of pCV-H77C). After washing, cells were incubated with FITC-conjugated goat anti-rabbit IgG (1:100) for 30 min at 37°C. After washing, slides were mounted and examined for immunofluorescence. For intracellular immunofluorescence staining, cells were fixed and permeabilized with cold acetone, as previously described (Forns et al., 1999).

Transfection of a chimpanzee with H77C(Δ HVR1) RNA

RNA was transcribed *in vitro* with T7 RNA polymerase from 10 μ g of template plasmid H77C(Δ HVR1) linearized with *Xba*I as described previously (Yanagi et al., 1997). The quality and amount of RNA were analyzed by gel electrophoresis and ethidium bromide staining. Two transcription mixtures were each diluted with 400 μ l of ice-cold phosphate-buffered saline without calcium or magnesium and then immediately frozen on dry ice and stored at -80°C. Within 24 hours, both transcription mixtures

were percutaneously injected into the liver of a chimpanzee (number 1590) under ultrasonographic control. The housing, maintenance, and care of the animals were in compliance with all relevant guidelines and requirements.

Serum samples were collected weekly from the chimpanzee and monitored for serum levels of alanine aminotransferase (ALT), anti-HCV antibodies [second generation ELISA] and HCV-RNA [HCV Monitor test (Roche) and in-house RT-nested PCR (Bukh et al., 1998)]. Sequence analysis of the recovered virus was performed at different time points during follow-up. In short, genomic regions were amplified in RT-nested PCR (Bukh et al., 1998) with primers specific for the H77 strain of HCV.

Construction of vectors that Express Mutant E2 Protein on the Cell Surface or in a Form That Is Secreted

A. Cell Surface Δ HVR1-E2 Vectors

Clones E1E2-715 (amino acids 192-715) and Δ HVR1-7 (amino acids 192-383 of E1 in frame with amino acids 411-715 of E2) were produced as described above.

Clone Δ HVR1-mut5 (amino acids 192-383 of E1 in frame with amino acids 411-715 of E2 except that amino acid 615 is changed from a leucine to a histidine) was produced as follows.

Primers E1-BglIII (SEQ ID NO: 4) and HVR1-mutR (SEQ ID NO: 8) (containing a T->A mutation at nucleotide position 2185: see Example 4) were used to amplify a fragment encoding the E1 protein and a portion of the E2 protein using Advantage KlenTaq Polymerase mix, as described above. Primers HVR-1 mutS (SEQ ID NO: 7) and E2-PstI (SEQ ID NO: 3) were used to amplify the rest of the truncated E2 protein by using AmpliTaq Gold DNA polymerase. In a second step, both products were fused by an

amplification reaction that contained a mixture of both external primers (E1-BglII (SEQ ID NO: 4) and E2-PstI (SEQ ID NO: 3) and the two fusion primers SEQ ID NO: 9 and SEQ ID NO: 10) (Table 1), by using Advantage KlenTaq Polymerase mix. The fusion product was digested with *BglII* and *PstI* and cloned into the digested pDisplay vector to produce ΔHVR1-mut5.

Clones E1E2-715, ΔHVR1-7 and ΔHVR1-mut5 were used to transfect Huh7 cells and to determine their pattern of reactivity against a panel of 5 rabbit hyperimmune sera and 12 human monoclonal antibodies (Cardoso *et al.*, 1998, Inchauspe *et al.* 1998) by immunofluorescence in live and fixed/permeabilized cells.

B. Secreted ΔHVR1-E2 Vectors

In addition, three different expression vectors encoding a secreted form of the E2 protein were constructed: E2-661, encoding the ER signal sequence of E1 (aa 364-383 of pCV-H77C) and a carboxy-terminal truncated E2 protein (aa 384-661 of pCV-H77C); E2-661 ΔHVR1, encoding the same protein with a deletion of the HVR1 region (aa 384-410 of pCV-H77C); and E2-661 ΔHVR1-mut, encoding the same protein with a deletion of the HVR1 and the replacement of leucine for histidine at amino acid position 615. In this case, amplification was performed from templates E1E2-715, ΔHVR1-7 and ΔHVR1-mut5, respectively, with primers shown in Table 1. PCR products were cloned into the expression vector pcDNA3.1(-) (Invitrogen). Clones E2-661, E2-661 ΔHVR1, and E2-661 ΔHVR1-mut were used to express soluble E2 protein *in vitro*. Briefly, one microgram of each of the plasmids was used for *in vitro* transcription-translation. Reactions were performed in 25 μl of the TNT Coupled Reticulocyte Lysate System (Promega) containing [³⁵S] methionine,

with the addition of canine microsomal membranes at 30°C for 90 min.

Example 1

In vitro transcription-translation analysis

To determine whether cleavage between E1 and E2 was affected by the deletion of the HVR1 region, expression vectors E1E2-715 (encoding the complete E1 protein and a cell-surface E2 protein lacking the carboxy-terminal 31 amino acids of E2) and ΔHVR1-7 (encoding the complete E1 protein and a cell-surface E2 lacking the HVR1 and carboxy-terminus 31 amino acids) were assayed for *in vitro* protein synthesis. *In vitro* transcription-translation was performed with the TNT Coupled Reticulolysate System using [S³⁵]methionine. In the absence of microsomal membranes, translation products of the expected size were obtained for E1E2-715 and ΔHVR1-7 (Figure 2). When microsomal membranes were added, cleavage between E1 and E2 was observed in both cases (Figure 2) and, as expected, the E2 protein derived from ΔHVR1-7 was slightly smaller in size. These results indicate that cleavage between E1 and E2 is not affected by the deletion of the N-terminus (HVR1) of the E2 protein.

Example 2

Analysis of cell-surface expression of the E2 protein

To analyze whether a truncated E2 protein lacking the HVR1 region could be targeted to the cell surface, Huh7 cells were transfected with E1E2-715 and ΔHVR1-7. After transfection, fixed/permeabilized cells and live cells were analyzed for expression of E2 by immunofluorescence. Up to 20% of cells transfected with E1E2-715 were positive when stained with a rabbit hyperimmune serum 1mf 86 or FOR-1. Both fixed

/permeabilized and live cells were stained. A similar proportion of cells transfected with Δ HVR1-7 was positive when stained with rabbit hyperimmune serum FOR-1 (raised against aa 517-535). Again, both fixed/permeabilized and live cells were stained. In contrast, there was no staining with rabbit hyperimmune serum lmf86, which was raised against aa 390-410 (within HVR1). These results demonstrated that transport of the E2 protein to the cell surface was not affected by deletion of the HVR1 region, suggesting that deletion of the N-terminal domain of E2 did not result in misfolding of this protein.

Example 3

Transfection of a chimpanzee with RNA of H77C(Δ HVR1)

To determine the infectivity of H77C(Δ HVR1), RNA transcripts were injected into the liver of chimpanzee 1590. HCV-RNA was detected by RT-nested PCR at week one post-inoculation (p.i.) and the animal remained HCV-RNA positive throughout the entire follow-up period of 22 weeks (Figure 3). The quantitative HCV Monitor test was negative until week 10 p.i. The genome titer during the first 7 weeks was \square 10 genome equivalent (GE)/ml. However, beginning at week 8, the titer progressively increased over time (Table 2 and Figure 3).

Table 2

Chimpanzee 1590 Infected with H77C(Δ HVR1)

Week	HCV RNA*	Titer† in house	Monitor‡	E2 (nt 2185)	NS3 (nt 3769)	NS5B (nt 8966)
0	-					
1	+/-					
2	+/-					
3	+	1				
4	+	1		T	G	
5	+	1		T	G	A
6	+	1		T	G	A
7	+	1	-	T	G	A
8	+	2	-	T/A	G	A/t
9	+	2	-	T/A	G/a	T/a
10	+	2	119	A/t	G/a	T/a
11	+	3	756	A	G/a	T
12	+	3	682	A	G/A	T
13	+	4	1981	A	A	T
14	+	4	3128			
15	+		3038			
16	+		3285			
17	+		1928			
18	+		5292			

* In-house RT-PCR qualitative test

† log₁₀ HCV GE titer by in-house RT-PCR‡ Amplicor HCV monitor (number of genome copies/ml) (Roche Diagnostics)
Nucleotide sequence corresponding to Yanagi 1997

Note: Lower case letters represent minor species

Serum liver enzyme values remained normal during the follow-up. The second generation anti-HCV test remained negative at least through week 22 p.i.

The follow-up period for chimpanzee 1590 has been extended to 72 weeks. Chimpanzee 1590 remained HCV-RNA positive throughout the 72 week follow-up period. Thus, the chimpanzee developed a chronic infection. In addition, the genome titer reached 10^3 - 10^4 GE/ml at weeks 11-14 (see Table 2) and remained at 10^3 - 10^4 GE/ml throughout the remainder of the 72 week period.

At weeks 51 and 52 p.i., the relative genome titer of HCV lacking HVR1 was also assessed in serum, PMBC, and liver obtained from the transfected chimpanzee. HCV RNA was not

detected in PBMCs, but the HCV titers determined from liver tissue at weeks 51 and 52 p.i. were higher than the titers determined from the serum.

The 72 week follow-up period also revealed that the transfected chimpanzee became positive for antibodies to second generation ELISA HCV at week 37 and third generation RIBA (Chiron) confirmed the presence of antibodies to C22(core), C33-c (NS3), and c100 (NS3-NS-4).

Finally, CD4+ and CD8+ T cell responses were also measured. An HCV-specific proliferative CD4+ T cell response to C22 (core), C33-c (NS3), c100(NS3-NS4) and/or NS5 antigens of HCV was detected in the peripheral blood mononuclear cells (PBMC) beginning at week one post-infection and throughout the follow-up period. During most weeks, however, only a mono-specific response to the core was detected.

Peripheral CD8+ T cell (CTL) responses were tested at weeks 5, 9, 14, and 18 by an *in vitro* peptide stimulation assay. CTL were detected with two NS4 peptides at weeks 9 and 14 and with one of these at week 18. A transient and monospecific (NS5) proliferative T cell response was detected in the liver at weeks 14 and 15. Although serum liver enzyme values remained normal during the entire follow-up, necroinflammatory changes indicative of hepatitis were detected in liver biopsies during weeks 34-40.

Example 4

Analysis of the HCV-RNA recovered from chimpanzee 1590

Sequence analysis of the HCV-RNA recovered from chimpanzee 1590 was performed at week 13. The entire ORF was amplified by RT-nested PCR of overlapping gene regions. Direct sequence analysis of both strands of DNA from the PCR products

confirmed that the recovered virus lacked the HVR1 region. Interestingly, three nucleotide mutations were identified; all three mutations were non-synonymous i.e. resulted in an amino acid change (Table 2). One mutation was located within the E2 region (corresponding to aa position 615 of H77C) and generated a change from leucine to histidine. Another mutation was located within the serine-protease domain of NS3 (corresponding to aa position 1143 of H77C; arginine to histidine). Both mutations are at positions that are universally conserved among all HCV genotype reference strains (Bukh et al., 1998). Finally, a third mutation was observed in the NS5B RNA-polymerase domain (corresponding to aa position 2875 of H77C; glutamic acid to aspartic acid). Although this is not a universally conserved position, a substitution of aspartic acid has not been described previously. Thus, all three mutations were unique.

Two additional mutations were identified upon rereading of the week 13 sequence: mutation at 514 (valine to methionine) and a synonymous mutation at amino acid 2631. Sequence analysis of the HCV-RNA recovered from chimpanzee 1590 was also performed at week 24 and five nucleotide mutations were identified; four of which were non-synonymous. (Table 3).

Table 3

Evolution of HCV lacking HVR1 in chimpanzee 1590*. We performed sequence analysis of the entire open reading frame of the virus recovered from chimpanzee 1590 at weeks 13 and 24 post-transfection. We confirmed that the HVR1 region was not present and identified differences from the wild-type cDNA clone as indicated.

Week	Titer In-house†	Titer Monitor‡	E2				NS3				NS5B			
			nt.1881	aa.514	nt.2185	aa.615	nt.3769	aa.1143	nt.4708	aa.1456	nt.8234	aa.2631	nt.896 6	aa.2875
			G	V	T	L	G	R	C	T	G	K	A	E
1	<1	Neg												
2	<1	Neg												
3	1	Neg												
4	1	Neg			T	L	G	R						
5	1	Neg	G	V	T	L	G	R					A	E
6	1	Neg	G	V	T	L	G	R					A	E
7	1	Neg	G	V	T	L	G	R					A	E
8	2	Neg	G	V	T/A	L/H	G	R					A/A	E/d
9	2	Neg	G	V	T/A	L/H	G/a	R/h					T/a	D/c
10	2	2.08	G	V	A/h	H/l	G/a	R/h					T/a	D/c
11	3	2.88	G	V	A	H	G/a	R/h					T	D
12	3	2.83	G/a	V/m	A	H	G/A	R/H					T	D
13	4	3.30	G/A	V/M	A	H	G/A	R/H	C	T	G/a	K	T	D
14	4	3.50	G/A	V/M			G/A	R/H	C	T	G	K		
15	3	3.48												
16	4	3.52	A/g	M/A			G/a	R/h	C	T	G/a	K		
17	4	3.29												
18	3	3.72	A	M			G	R	C/h	T/m	G/A	K		
19	3	3.20												
20	3	3.37	A	M			G	R	C/T	T/M	A/g	K		
21	3	4.04												
22	4	3.94	A	M			G	R	C/T	T/M	A/g	K		
23	4	3.48												
24	4	3.74	A	M	A	H	G	R	T/c	M/l	A	K	T	D

*Nucleotide (nt) and amino acid (aa) positions of H77C. Sequence of H77C(ΔHVR1) cDNA is shown on top. Dominant sequences recovered from the chimpanzee are shown in capital letters. Minor sequences recovered from the chimpanzee are shown in lower case letters. †log₁₀ titer determined by RT-nested PCR on 10-fold serially diluted extracted RNA. ‡log₁₀ titer determined by second generation Monitor test (Roche).

Four of the mutations were identical to the mutations identified at week 13 and one of the week 13 mutations (the NS3 mutation at amino acid 1143) reverted to wild-type. The single new mutation identified at week 24 was located within the serine protease domain of NS3 (corresponding to aa position 1456 of H77C; threonine to methionine).

To analyze a possible relationship between the increase in viral titer and the appearance of the four non-synonymous mutations identified at week 13, sequence analysis was performed from regions encompassing the described mutations. During weeks 4-7, direct sequence analysis did not reveal evidence of the mutated nucleotides at the four positions. Between weeks 8 and 14, coinciding with the first increase in viral titer, direct sequence analysis revealed the presence of amino acid substitutions at four positions (two in E2, one in NS3, and one in NS5B) (See Table 3). Therefore, there was a temporal association between the appearance of these mutations and an increase in viral titer (See Tables 2 and 3).

Example 5

Transmission of HCV lacking HVR1 to a second chimpanzee

To investigate whether the HCV mutant lacking HVR1 was transmissible, a naïve chimpanzee (#96A008) was inoculated intravenously with 90 ml of plasma taken at week 4 from chimpanzee 1590, before appearance of consensus mutations in that chimpanzee (Table 3). The virological and immunological correlates of HCV infection in chimpanzee 96A009 are shown in Fig. 4. Serum HCV-RNA was first detected at week 2 p.i., with a genome titer of 10^1 GE/ml and the titer increased to peak levels of 10^4 - 10^5 GE/ml during weeks 3-9. The infection was resolved at week 18 p.i. Antibodies to HCV were not detected. A peripheral

mono-specific proliferative CD4+ T cell response against core (C22) was detected during weeks 4-13 p.i.; a multi-specific response against NS3 (C33-c), NS3-NS4 (c100) and NS5 was detected at week 18. The chimpanzee mounted an early and multi-specific peripheral CD8+ T cell response to a total of seven different epitopes (representing core, NS3, NS4 and NS5) at the weeks tested (weeks 4, 7, 11, 15 and 19). Finally, a multi-specific (C33-c, c100 and NS5) sustained proliferative CD4+ T cell response was detected in the liver from week 8 p.i. Liver enzyme values were marginally elevated during weeks 10-18 p.i. and necroinflammatory changes were detected in liver biopsies during weeks 11-19 p.i.

Sequence analysis of the entire open reading frame of HCV recovered from the chimpanzee 96A008 at weeks 4 and 9 p.i. showed that the transmitted virus lacked HVR1. Compared with the cDNA clone of that mutant, four nucleotide substitutions were identified in the virus and all four mutations were non-synonymous. (Table 4).

Table 4

Evolution of HCV lacking HVR1 in chimpanzee 96A008*. We performed sequence analysis of the entire open reading frame of the virus recovered from chimpanzee 96A008 at weeks 4 and 9 post-inoculation. We confirmed that the HVR1 region was not present and identified differences from the wild-type cDNA clone as indicated.

Week	Titer In-house†	Titer Monitor‡	E1		E2		NS3			
			nt.1097	aa.252	nt.1881	aa.514	nt.3769	aa.1143	nt.4269	aa.1310
			A	K	G	V	G	R	T	Y
1	Neg	Neg								
2	1	Neg	A/T	K/N	A	M	G	R	C	H
3	3	3.33	T	N	A	M	G/a	R/h	C	H
4	4	4.00	T	N	A	M	G/a	R/h	C	H
5	4	3.98					G/A	R/H		
6	5	4.25					G/A	R/H		
7	5	4.81					G/A	R/H		
8	5	4.51					G/A	R/H		
9	5	4.62	T	N	A	M	G/A	R/H	C	H
10	4	3.30					G/a	R/h		
11	3	2.63								
12	3	2.92					G	R		
13	2	Neg								
14	2	3.14					G	R		
15	1	Neg								
16	4	3.06					G	R		
17	1	Neg								
18	Neg	Neg								
19	Neg	Neg								
20	Neg	Neg								

*Nucleotide (nt) and amino acid (aa) positions of H77C. Sequence of H77C(ΔHVR1) cDNA is shown on top. Dominant sequences recovered from the chimpanzee are shown in capital letters. Minor sequences recovered from the chimpanzee are shown in lower case letters. †log₁₀ titer determined by RT-nested PCR on 10-fold serially diluted extracted RNA. ‡log₁₀ titer determined by second generation Monitor test (Roche).

Sequence analysis of regions encompassing the described mutations showed that a mutation in E2 and NS3 each had replaced the wild-type nucleotide by the time the virus was first detected (week 2). An identical change in E2 had occurred in the transfected chimpanzee (Table 3). A change in E1 appeared as the quasispecies at week 2 but had replaced the wild-type nucleotide completely by week 3. The other mutation, in NS3, appeared at week 3 and persisted as a quasispecies during weeks 3-10 and then reverted to wild-type. This latter mutation had been seen also in the transfected chimpanzee. (Table 3).

Example 6Analysis of potential conformational changes
within E2 after L615H substitution

Sequence analysis of HCV recovered from chimpanzee 1590 demonstrated that a non-synonymous mutation occurred within the E2 protein during follow-up (nt 2186 of pCV-H77C; amino acid position 615). Since a change from a non-polar amino acid (leucine) to a charged one (histidine) within E2 might change its conformation, Huh7 cells were transfected with E1E2-715 (intact E1-E2), Δ HVR1-7 (the deletion of HVR1) or Δ HVR1-mut (deletion of HVR1 along with the mutation at position 615), and the transfected cells were stained for E2 with a panel of 4 rabbit hyperimmune sera and 12 human monoclonal antibodies (Cardoso et al. 1998; Inschauspe et al., 1998). Immunofluorescence analysis showed that there was no significant change in the patterns of reactivity among the three different forms of E2 targeted to the cell surface (Table 5).

Table 5

Effect of deletion of the HVRI region and mutation of
E2 615 (leu-his) on the reactivity of cell surface-expressed E2

<u>Antibody</u>	<u>E1E2-715</u>	<u>ΔHVRI-7</u>	<u>ΔHVRI-mut5</u>
RHS 1mf86 (aa390-410)	++++	-	-
RHS 521 (aa 645-662)	++	+++	++
RHS FOR-1(aa 517-535)	+++	+++	+++
RHS 554 (aa 460-483)	++++	+++	+++
2H1	+/-	-	-
2G1	++	++	+
1H3	++	+++	+
4F7	++	++	+
8E8	-	-	-
2H8	+++	+++	++
4E5	++	+++	++
4F1	++	+++	+++
7B7	+++	++	++
108F6	++	+	++
503E7p	+++	++	+
29E4p	++	++	+

-RHS: rabbit hyperimmune sera.

-Amino acid position within the H77C sequence of the peptides used to generate RHS are indicated in parenthesis.

-The remaining 12 antibodies are human monoclonal antibodies (Cardoso et al. 1998; Inschauspe et al., 1998).

-Scoring system from - to +++. - means no staining and +++ means strong staining.

References

- Alter, M. J. (1997). Hepatology **26**, 62S-65S.
- Bukh, J., Miller, R.H., and Purcell, R.H. (1995) *Semin. Liver Dis.* **15**, 41-63
- Bukh, J., et al. (1998) *Hepatology* **28**, 319A
- Bukh, J., et al. (1998) *J. Infect. Dis.* **178**:1193-1197
- Cardoso, M., et al., (1998) *J. Med. Virol.* **55**:28-34
- Davis, G. L., et al., *N. Engl. J. Med.* **339**, 1493-1499.
- Farci, P., et al. (1996) *Proc. Natl. Acad. Sci. USA* **93**, 15394-15399
- Forns, X., and Bukh, J. (1998) *Viral Hepatitis Rev.* **4**, 1-19
- Hijikata, M., et al. (1991) *Biochem Biophys. Res. Commun.* **175**, 220-228
- Hoofnagle, J. H. (1997). *Hepatology* **26**, 15S-20S.
- Harbersetzer, F., et al. (1998) *Virology* **249**:32-41
- McAllister, J., et al. (1998) *J. Virol.* **72**: 4893-4905
- McHutchison, J. G., et al. (1998) *N. Engl. J. Med.* **339**, 1485-1492.
- Puntoriero, G., et al. (1998) *EMBO J.* **17**:3521-3533
- Ray, S.C., et al. (1999) *J. Virol.* **73**, 2938-2946
- Rice, C.M. (1996) in *Fields Virology* (Fields, B.N., Knipe, D.M. and Howley, P.M. eds), pp 931-960, Lippincott-Raven
- Shimizu, Y., et al. (1996) *Virology* **223**:409-414
- Taylor, D., et al. (1999) *Science* **285**:107-110
- Weiner, A.J., et al. (1991) *Virology* **180**, 842-848
- Weiner, A.J., et al. (1992) *Proc. Natl. Acad. Sci. USA* **89**, 3468-3472
- Yanagi, M., et al. (1997) *Proc. Natl. Acad. Sci. USA* **94**, 8738-8743
- Yanagi, M., et al. (1999) *Virology* **262**, 250-263
- Zibert, A., et al. (1997) *Hepatology* **25**, 1245-1249

CLAIMS

1. A nucleic acid molecule which encodes a human hepatitis C virus lacking the hypervariable region 1 of HCV envelope 2 protein, said molecule capable of expressing said virus when transfected into cells.

2. The nucleic acid molecule of claim 1, wherein the molecule encodes at least one of the following amino-acids, a histidine at amino acid positions 615, 1143 and 1310, an aspartic acid at position 2875, a methionine at amino acid positions 514 and 1456, and an asparagine at amino acid position 252.

3. A nucleic acid molecule comprising a chimeric virus genome, said genome being a pestivirus or flavivirus genome in which the structural region of the pestivirus or flavivirus has been replaced by the structural region of a hepatitis C virus genome which lacks hypervariable region 1 of HCV envelope 2 protein.

4. The nucleic acid molecule of claim 3, wherein at least one envelope gene from the structural region of the pestivirus or flavivirus genome has been replaced by the E2 gene from the structural region of a hepatitis C virus genome, said HCV E2 gene lacking hypervariable region one.

5. The nucleic acid molecule of claim 3, wherein the structural region of the hepatitis C virus genome encodes at least one of the following amino-acids, a histidine at amino acid position 615, an asparagine at amino acid position 252, and a methionine at position 514.

6. The nucleic acid molecule of claims 3-5, wherein the genome is a pestivirus genome.

7. The nucleic acid molecule of claims 3-5, wherein the pestivirus genome is a bovine viral diarrhea virus genome.

8. The nucleic acid molecule of claims 3-5, wherein the genome is a flavivirus genome.

9. The nucleic acid molecule of claim 8, wherein the flavivirus genome is the genome of a dengue virus.

10. A DNA construct comprising a nucleic acid molecule according to claim 1.

11. A DNA construct comprising a nucleic acid molecule according to claim 3.

12. An RNA transcript of the DNA construct of claim 10.

13. An RNA transcript of the DNA construct of claim 11.

14. A host cell transformed or transfected with the DNA construct of claim 10.

15. A host cell transformed or transfected with the DNA construct of claim 11.

16. A host cell transformed or transfected with RNA transcript of claim 12.

17. A host cell transformed or transfected with RNA transcript of claim 13.

18. A hepatitis C virus envelope 2 polypeptide produced by the cell of claims 14, 15, 16 or 17.

19. A hepatitis C virus produced by the cell of claims 14 or 16.

20. A chimeric virus produced by the cell of claims 15 or 17.

21. A hepatitis C virus whose genome comprises a nucleic acid molecule according to claim 1.
22. A chimeric virus whose genome comprises a nucleic acid molecule according to claim 3.
23. A host cell infected with the virus of claim 21.
24. A host cell infected with the virus of claim 22.
25. An envelope 2 polypeptide encoded by the nucleic acid sequence of claims 1 or 3.
26. Antibody to the polypeptide of claim 25.
27. Antibody to the hepatitis C virus of claim 21.
28. Antibody to the chimeric virus of claim 22.
29. A composition comprising a polypeptide of claim 25 suspended in a suitable amount of a pharmaceutically acceptable diluent or excipient.
30. A composition comprising a virus of claims 21 or 22 suspended in a suitable amount of a pharmaceutically acceptable diluent or excipient.
31. A composition comprising a nucleic acid molecule of claims 1 or 3 suspended in a suitable amount of a pharmaceutically acceptable diluent or excipient.
32. A method of immunizing a mammal, said method comprising immunizing said mammal with the composition of claim 29.
33. A method of immunizing a mammal, said method comprising immunizing said mammal with the composition of claim 30.

34. A method of immunizing a mammal, said method comprising immunizing said mammal with the composition of claim 31.
35. Antibodies produced by the method of claim 32.
36. Antibodies produced by the method of claim 33.
37. Antibodies produced by the method of claim 34.
38. T cells reactive with the virus of claims 21 or 22, said T cells produced by the method of claim 33.
39. T cells reactive with the virus of claims 21 or 22, said T cells produced by the method of claim 34.
40. A chimeric gene comprising, in 5' to 3' order:
(i) an endoplasmic reticulum signal sequence;
and
(ii) a coding sequence which encodes a hepatitis C virus (HCV) envelope 2 protein lacking a)
hypervariable region 1 of the full-length
envelope 2 protein and b) at least the 30
carboxy-terminal amino acids of the full-length
envelope 2 protein.
41. The chimeric gene of claim 40, wherein the envelope 2 protein coding sequence lacks about the carboxy terminal 85 amino acids of the full-length envelope 2 protein.
42. The chimeric gene of claim 40, wherein the endoplasmic reticulum signal sequence of i) is amino acids 364-384 of the HCV E1 gene.
43. The chimeric gene of claim 40, wherein the gene further comprises a plasma membrane anchor sequence 3' of the envelope 2 protein coding sequence of ii).

44. The chimeric gene of claim 43, wherein an E1 coding sequence is located between the ER signal sequence of i) and the envelope 2 coding sequence of ii).

45. An expression vector comprising the chimeric gene of claim 40.

46. An expression vector comprising the chimeric gene of claim 43.

47. A host cell transformed or transfected with the expression vector of claim 45.

48. A host cell transformed or transfected with the expression vector of claim 46.

49. A method for expressing a secreted hepatitis C virus envelope 2 protein lacking hypervariable region, said method comprising transforming a host cell with the expression vector of claim 45 under conditions which permit expression of the secreted envelope 2 protein.

50. A method for expressing a hepatitis C virus envelope 2 protein lacking HVR1 on the surface of a cell, said method comprising transforming a host cell with the expression vector of claim 46 under conditions which permit expression of the envelope 2 protein on the cell surface.

51. An envelope 2 protein produced by the method of claim 49.

52. An envelope 2 protein produced by the method of claim 50.

53. A method for immunizing a mammal comprising administering the expression vector of claim 45 to the mammal in an amount effective to stimulate the production of protective antibodies to HCV.

54. A method for immunizing a mammal comprising administering the expression vector of claim 46 to the mammal in an amount effective to stimulate the production of protective antibodies to HCV.

55. A method for immunizing a mammal comprising administering the envelope 2 protein of claim 51 to the mammal in an amount effective to stimulate the production of protective antibodies to HCV.

56. Antibodies produced by the method of claims 53, 54 or 55.

57. A pharmaceutical composition comprising the antibodies of claim 56.

58. A pharmaceutical composition comprising the expression vector of claims 45 or 46.

59. A pharmaceutical composition comprising the protein of claim 51.

60. The nucleic acid molecule of claim 2, wherein the molecule encodes a methionine at amino acid position 514.

FIG. 1A

10	20	30	40	50	
1234567890	1234567890	1234567890	1234567890	1234567890	
GCCAGCCCC	TGATGGGGGC	GACACTCCAC	CATGAATCAC	TCCCTGTGA	50
GGAACTACTG	TCTTCACGCA	GAAAGOGTCT	AGCCATGGCG	TTAGTATGAG	100
TGTGTGTCAG	CCTCCAGGAC	CCCCCTCCC	GGGAGAGCCA	TAGTGGTCTG	150
CGGAACOGGT	GAGTACACCG	GAATTGCCAG	GACGACOGGG	TCCTTTCTTG	200
GATAAACCCG	CTCAATGCCT	GGAGATTTGG	GOGTGGCCCC	GCAAGACTGC	250
TAGCCGAGTA	GIGTGGGGTC	GCGAAAGGCC	TTGTGGTACT	GCCTGATAGG	300
GTCCTTGGCA	GIGCCCCGGG	AGGTCTCGTA	GACCGTGCAC	CATGAGCACG	350
AATCCTAAAC	CTCAAAGAAA	AACCAAACGT	AACACCAACC	GTGCCCCACA	400
GGACGTCAG	TTCCCCGGGTG	GCGGTCAGAT	CGTTGGTGGG	GTTTACTTGT	450
TGCGCGCAG	GGGCCCTAGA	TTGGGTGTGC	GCGCGACGAG	GAAGACTTCC	500
GAGCGGTGCG	AACCTCGAGG	TAGAOGTCAG	CCTATCCCCA	AGGCACGTCC	550
GCCCCAGGGC	AGGACCTGGG	CTCAGCCCCG	GTACCCCTTG	CCCCCTCTATG	600
GCAATGAGGG	TTGCGGGTGG	GCGGGATGGC	TCCTGTCTCC	CCGTGGCTCT	650
CGGCTAGCT	GGGGCCCCAC	AGACCCCCGG	CGTAGGTCCG	GCAATTTGGG	700
TAAGGTCATC	GATACCCCTA	CGTGCGGCTT	CGCCGACCTC	ATGGGGTACA	750
TACCGCTCGT	CGCGCCCCCT	CTTGGAGGGG	CTGCCAGGGC	CCTGGCGCAT	800
GGGTCCGGG	TTCTGGGAGA	CGGCGTGAAC	TATGCAACAG	GGAACTTCC	850
TGGTTGCTCT	TTCTCTATCT	TCCTTCTGGC	CCCTGCTCTT	TGCTGACTTG	900
TGCCCCCTTC	AGCCTACCAA	GTGCGCAATT	CCTCGGGGCT	TTACCATGTC	950
ACCAATGATT	GCCCTAACTC	GAGTATTGIG	TACGAGGCGG	CCGATGOCAT	1000
CCTGCACACT	CCGGGGTGTG	TCCCTTGGGT	TGGCGAGGGT	AACGCTCGA	1050
GGTGTGGGT	GGGGGTGACC	CCACGGTGG	CCACCAGGGA	CGGCAAATC	1100
CCACAACGC	AGCTTCGACG	TCATATOGAT	CTGCTGTGCG	GGAGCGGCAC	1150
CCTCTGCTCG	GCCCTCTACG	TGGGGACCT	GTCGGGTCT	GTCTTTCTTG	1200
TTGGTCAACT	GTTTACCTTC	TCTCCAGGC	GCCACTGGAC	GACGCAAGAC	1250
TGCAATTGTT	CTATCTATCC	CGGOCATATA	ACGGGTATC	GCATGGCATG	1300
GGATATGATG	ATGAACCTGG	CCCCACGGC	AGGTTGGTGG	GTAGCTCAGC	1350
TGCTCCGGAT	CCACAAGGC	ATCATGGACA	TGATCGCTGG	TGCTCACTGG	1400
GGAGTCCCTG	CGGGCATAGC	GTATTTCTCC	ATGGTGGGGA	ACTGGGCGAA	1450
GGTCCCTGGTA	GTCCTGCTGC	TATTTGCGGG	CGTCGACGGG	ATCCAACTGA	1500
TCAACACCAA	CGGCAGTTGG	CACATCAATA	GCACGGCCTT	GAATTGCAAT	1550
GAAAGCCTTA	ACACCGGCTG	GTTAGCAGGG	CTCTTCTATC	AACACAAATT	1600
CAACTCTTCA	GGCTGTCCCTG	AGAGGTGGC	CAGCTGCCGA	CGCCTTACCG	1650
ATTTTGGCCA	GGGCTGGGGT	CCTATCAGTT	ATGCCAACGG	AAGCGGCCTC	1700
GACGAACGCC	CCTACTGCTG	GCACTACCCCT	CCAAGACCTT	GTGGCATTGT	1750
GCCCCCAAAG	AGGTTGTGTG	GCCCCGATATA	TTGCTTCACT	CCAGCCCCCG	1800

FIG. 1B

10	20	30	40	50	
1234567890	1234567890	1234567890	1234567890	1234567890	
TGGTGGTGGG	AACGACCGAC	AGGTGGGGCG	CGCCTAOCCTA	CAGCTGGGGT	1850
GCAAATGATA	CGGATGICIT	CGTCTTAAC	AACACCAGGC	CACCGCTGGG	1900
CAATTGGTTC	GGTTGTACCT	GGATGAACTC	AACTGGATTG	ACCAAAGTGT	1950
GCGGAGCGCC	CCCTTGITGIC	ATCGGAGGGG	TGGGCAACAA	CACCTTGCTC	2000
TGCCCCACTG	ATTGCTTCCG	CAAACATCCG	GAAGCCACAT	ACTCTGGGTG	2050
CGGCTCCGGT	CCCTGGATTG	CACCCAGGIG	CATGGTGGAC	TACCCGTATA	2100
GGCTTTGGCA	CTATCTTGT	AACATCAATT	ACACCATATT	CAAAGTCAGG	2150
ATGTACGTGG	GAGGGGTGGA	GCACAGGCTG	GAAGCGGCTT	GCAACTGGAC	2200
GCGGGGGGAA	CGCTGTGATC	TGGAAGACAG	GGACAGGTCG	GAGCTCAGCC	2250
CGTGTCTGCT	GTCCACCACA	CAGTGGCAGG	TCTTCCGGTG	TCTTTTCAAG	2300
AACCTGOCAG	CCTTGITCCAC	CGGCTCATC	CACCTCCACC	AGAACATTGT	2350
GGACGTGCAG	TACTTGTACG	GGGTAGGGTC	AAGCATGGCG	TCTTGGGCGA	2400
TTAAGTGGGA	GTACGTGCTT	CTCTGTTCG	TTCTGCTTGC	AGACGGCGGC	2450
GTCTGCTCCT	GCTTGITGGAT	GATGTTACTC	ATATCCCAAG	CGGAGCGGCG	2500
TTTGGAGAAC	CTCGTAATAC	TCAATGCAGC	ATCCCTGGCC	GGGAGCGACG	2550
GTCTGTGTGC	CTTCTCTGTG	TTCTTCTGCT	TTGGGTGGTA	TCCTAAGGGT	2600
AGGTGGGTGC	CCGGAGCGGT	CTACGCCCTC	TACGGGATGT	GGCCTCTCCT	2650
CCTGCTCCTG	CTGGCGTTGC	CTCAGCGGGC	ATACGCACTG	GACAGGGAGG	2700
TGGCGCGGTC	GTGTGGCGGC	GTGTGTTCTG	TGGGGTTAAT	GGGCTGACT	2750
CTGTGCGCAT	ATTACAAGCG	CTATATCAGC	TGGTGCATGT	GGTGGCTTCA	2800
GTATTTTCTG	ACCAGAGTAG	AAGCGCAACT	GCACGTGTGG	GTCCCCCCCC	2850
TCAACGTCCG	GGGGGGGGCG	GATGCGGTCA	TCTTACTCAT	GTGTGTAGTA	2900
CACCCGACCC	TGGTATTTGA	CATCACCAAA	CTACTCCTGG	CCATCTTCCG	2950
AACCCCTTGG	ATTCTTCAAG	CCAGTTTGGT	TAAAGTCCCC	TACTTCTGTC	3000
GCGTCAAGG	CCCTCTCCGG	ATCTGGCGCG	TAGCGCGGAA	GATAGCCGGA	3050
GGTCATTACG	TGCAAATGGC	CATCATCAAG	TTAGGGGGCG	TTACTGGCAC	3100
CTATGIGTAT	AACCATCTCA	CCCTCTTCCG	AGACTGGGCG	CACAACGGCC	3150
TGCGAGATCT	GGCGGTGGCT	GTTGTAACAG	TGCTCTTCTC	CCGAATGGAG	3200
ACCAAGCTCA	TCAAGTGGGG	GGCAGATACC	GCCGGGTGGG	GTGACATCAT	3250
CAACGGCTTG	CCGCTCTCTG	CCGTAGGGGG	CCAGGAGATA	CTGCTTGGGC	3300
CAGCCGACGG	AATGGTCTCC	AAGGGGTGGG	GGTTGCTGGC	GGCCATCACG	3350
GCGTACGCCC	AGCAGACGAG	AGGCTCTCTA	GGGTGTATAA	TCAACAGCCT	3400
GACTGGCGCG	GACAAAAACC	AAGTGGAGGG	TGAGGTCCAG	ATCGTGTCAA	3450
CTGCTACCCA	AACCTTCTCTG	GCAACGTGCA	TCAATGGGGT	ATGCTGGACT	3500
GTCTACCCAG	GGGCGGGAAC	GAGGACCATC	GCATCACCCA	AGGGTCTCTG	3550
CATCCAGATG	TATACCAATG	TGGACCAAGA	CCTTGTGGGC	TGGCCCGCTC	3600

FIG. 1C

10	20	30	40	50	
1234567890	1234567890	1234567890	1234567890	1234567890	
CTCAAGGTC	CCGCTCATTG	ACACCTGTGA	CCTGCGGCTC	CTCGGACCTT	3650
TACCTGGTCA	CGAGGCAAGC	CGATGTCATT	CCCGTGGGCC	GGCGAGGTGA	3700
TAGCAGGGGT	AGCCTGCTTT	CGCCCCGGCC	CATTTCCCTAC	TTGAAAGGCT	3750
CCTCGGGGGG	TCCGCTGTTG	TGCCCCGGCG	GACACGGCGT	GGGCTATATC	3800
AGGGCCGGCG	TGTGCACCCG	TGGAGTGGCT	AAAGCGGTGG	ACTTTATCCC	3850
TGTGGAGAAC	CTAGGGACAA	CCATGAGATC	CCCGTGTGTC	AOGGACAACT	3900
CCTCTCCACC	AGCAGTGGCC	CAGAGCTTCC	AGGTGGCCCA	CCTGCATGCT	3950
CCCACCGGCA	GCGGTAAAGAG	CACCAAGGTC	CCGGCTGGGT	ACGCAGCCCA	4000
GGGCTACAAG	GTTGTGGTGC	TCAACCCCTC	TGTTGCTGCA	ACGCTGGGCT	4050
TTGGTGCTTA	CATGTCCAAG	GCCATGGGG	TTGATCCCTAA	TATCAGGACC	4100
GGGGTGAGAA	CAATTACCAC	TGGCAGCCCC	ATCAGTFACT	CCACCTACGG	4150
CAAGTTCCCT	GCCGACGGCG	GGTGCCTCAG	AGGTGCTTAT	GACATAATAA	4200
TTTGTGACGA	GTGCCACTCC	ACGGATGCCA	CATCCATCTT	GGGCATUGGC	4250
ACTGTCTTIG	AOCAAGCAGA	GACTGGGGGG	GCGAGACTGG	TTGTGCTGGC	4300
CACTGCTACC	CCTCGGGGCT	CCGTCACTGT	GTCCCATCCT	AACATCGAGG	4350
AGGTTGCTCT	GTCCACCACC	GGAGAGATCC	CCTTTTAAGG	CAAGGCTATC	4400
CCCTCGAGG	TGATCAAGGG	GGGAAGACAT	CTCATCTTCT	GCCACTCAAA	4450
GAAGAAGTGC	GAAGAGCTCG	CCGGAAGCT	GGTGGCATTG	GGCATCAATG	4500
CCGTGGCCTA	CTACCGGGGT	CTTGACGTGT	CTGTCAATCC	GACCAGCGGC	4550
GATGTTGTGG	TGGTGTGGAC	CGATGCTCTC	ATGACTGGCT	TTACCGGGCA	4600
CTTCGACTCT	GTGATAGACT	GCAACAGTGT	TGTCACTCAG	ACAGTCGATT	4650
TCAGCCTTGA	CCCTACCTTT	ACCATTGAGA	CAACCAAGCT	CCCCCAGGAT	4700
GCTGTCTCCA	GGACTCAACG	CCGGGGCAGG	ACTGGCAGGG	GGAAGCCAGG	4750
CATCTATAGA	TTTGTGGCAC	CGGGGGAGCG	CCCTCGGGC	ATGTTGGACT	4800
CGTCGGTCT	CTGTGAGTGC	TATGACGGCG	GCTGTGCTTG	GTATGAGCTC	4850
ACGCCCCCGG	AGACTACAGT	TAGGCTACGA	GCGTACATGA	ACACCCCGGG	4900
GCTTCCCGTG	TGCCAGGACC	ATCTTGAATT	TTGGGAGGGC	GTCTTTACGG	4950
GCCTCACTCA	TATAGATGCC	CACTTTTTAT	CCAGACAAA	GCAGAGTGGG	5000
GAGAAGTTTC	CTTACCTGGT	AGCGTACCAA	GCCACCGTGT	GCGCTAGGGC	5050
TCAAGCCCCCT	CCCCCATCGT	GGGACCAGAT	GTTGAAGTGT	TTGATCGGCC	5100
TTAAACCCAC	CCTCCATGGG	CCAACACCCC	TGCTATACAG	ACTGGGGGCT	5150
GTTCAGAATG	AAGTCACCTT	GACGCACCCA	ATCACCAAT	ACATCATGAC	5200
ATGCATGTGG	GCGACCTGG	AGGTGCTCAC	GAGCACCTGG	GTGCTGTTTG	5250
GCGGGTCTCT	GGCTGCTCTG	GCCGGTATTT	GCTGTCTAAC	AGGCTGGGTG	5300
GTCATAGTGG	GCAGGATCGT	CTTGTCCGGG	AAGCCGGCAA	TTATACCTGA	5350
CAGGGAGGTT	CTCTACCAGG	AGTTGGATGA	GATGGAAGAG	TGCTCTCAGC	5400

FIG. 1D

10	20	30	40	50	
1234567890	1234567890	1234567890	1234567890	1234567890	
ACTTACCGTA	CATCGAGCAA	GGGATGATGC	TGCTGAGCA	GTTCAGCAG	5450
AAGGCCCTCG	GCTTCCTGCA	GACCGCGTCC	CGCATGCAG	AGGTATAC	5500
COCTGCTGTC	CAGACCAACT	GGCAGAACT	CGAGGCTTT	TGGGCGAAGC	5550
ACATGTGGAA	TTTCATCAGT	GGGATACAAT	ACTTGGCGGG	CCGTGCAACG	5600
CTGCTGGTA	AACCCGCCAT	TGCTTCATTC	ATGGCTTTTA	CAGCTGCGGT	5650
CACCAGCCCA	CTAACCACTG	GCCAAACCT	CTCTTCAAC	ATATTGGGGG	5700
GGTGGGTGGC	TGCCCAGCTC	GCGCCCCCG	GTCGCGCTAC	TGCTTTGTTG	5750
GGTGCTGGCC	TAGCTGGCGC	CGCCATCGGC	AGCGTTGGAC	TGGGGAAGGT	5800
OCTCGTGGAC	ATTCTTGCAG	GGTATGGCGC	GGCGTGGCG	GGAGCTCTTG	5850
TAGCATTCAA	GATCATGAGC	GGTGAGGTCC	CTCCACCGA	GGACCTGGTC	5900
AATCTGCTGC	CCGCCATCCT	CTCGCCCTGA	GCCCCTGTAG	TGGGTGTGGT	5950
CTGCGCAGCA	ATACTGCGCC	GGCAGGTTCG	CCCGCGCGAG	GGGGCAGTCC	6000
AATGGATGAA	CCGGCTAATA	GCCTTGGCCT	CCCGGGGGAA	CCATGTTTCC	6050
CCACGCACT	ACGTGCGCGA	GAGCGATGCA	GCCGCCCGCG	TCACGTCCAT	6100
ACTCAGCAGC	CTCAGTGTA	CCAGCTCCT	GAGGCGACTG	CATCAGTGA	6150
TAAGCTCGGA	GTGTACCACT	CCATGCTCCG	GTCTCTGGCT	AAGGGACATC	6200
TGGGACTGGA	TATGCGAGGT	GCTGAGCGAC	TTTAAGACCT	GGCTGAAAGC	6250
CAAGCTCATG	CCACAACCTG	CTGGGATTCC	CTTTGTGTCC	TGCCAGCGCG	6300
GGTATAGGGG	GGTCTGGCGA	GGAGACGGCA	TTATGCACAC	TGCTGCCAC	6350
TGTGGAGCTG	AGATCACTGG	ACATGTCAAA	AACGGGACGA	TGAGGATCGT	6400
CGGTCCTAGG	ACCTGCAGGA	ACATGTGGAG	TGGGACGTTC	CCCATTAACG	6450
OCTACACCAC	GGGCCCCGTG	ACTCCCTTTC	CTGCGCCGAA	CTATAAGTTC	6500
GCGCTGTGGA	GGGTGTCTGC	AGAGGAATAC	GTGGAGATAA	GGCGGGTGGG	6550
GGACTTCCAC	TACGTATCGG	GTATGACTAC	TGACAATCTT	AAATGCCCGT	6600
GCCAGATCCC	ATGCCCCGAA	TTTTTCACAG	AATTGGACGG	GGTGGCCCTA	6650
CACAGGTTTG	CGCCCCCTTG	CAAGCCCTTG	CTGCGGGAGG	AGGTATCATT	6700
CAGAGTAGGA	CTCCAGAGT	ACCCGGTGGG	GTGCAATTA	CTTGGCGAGC	6750
CCGAACCGGA	CGTAGCCGTG	TTGACGTCOA	TGCTCACTGA	TCCCTCCCAT	6800
ATAACAGCAG	AGGCGGCCGG	GAGAAGGTTG	GCGAGAGGGT	CACCCCTTTC	6850
TATGGCCAGC	TCTTGGCTA	GCCAGCTGTC	CGCTCCATCT	CTCAAGGCAA	6900
CTTGCAACGC	CAACCATGAC	TCCCTTGACG	CCGAGCTCAT	AGAGGCTAAC	6950
CTCTGTGGTA	GGCAGGAGAT	GGCGGGCAAC	ATCAACAGGG	TTGAGTCAGA	7000
GAACAAAGTG	GTGATCTTGG	ACTCCTTCGA	TCGCTTTGTG	GCAGAGGAGG	7050
ATGAGCGGGA	GGTCTCCGTA	CCTGCAGAAA	TTCTGCGGAA	GTCTCGGAGA	7100
TTGCCCCGGG	CCCTGCCCCG	CTGGGCGCGG	CCGACTACA	ACCCCCCGCT	7150
AGTAGAGACG	TGGAAAAAGC	CTGACTACGA	ACCACCTGTG	GTCCATGGCT	7200

FIG. 1E

10	20	30	40	50	
1234567890	1234567890	1234567890	1234567890	1234567890	
G000GCTACC	ACCTCCACGG	T000CT0CTG	TG0CT00G0C	TCGGAAAAAG	7250
CGTA0GGTGG	TCCTCACCGA	ATCAAC0CTA	TCTACTG0CT	TGG0CGAGCT	7300
TG0CACCAAA	AGTTTTGGCA	GCT0CTCAAC	TT00GGCATT	ACGGGCGACA	7350
ATA0GACAAC	AT0CTCTGAG	000G0000CT	CTGGCTG00C	000CGACT0C	7400
GACGTTGAGT	CCTATTCTTC	CATG00000C	CTGGAGGGGG	AG0CTGGGGG	7450
T00GGATCTC	AGCGACGGGT	CATGGT0GAC	GGTCAGTAGT	GGGG0CGACA	7500
0GGAAGATGT	CGTGTGCTGC	TCAATGTCTT	ATT0CTGGAC	AGGCGCACTC	7550
GTCAC000GT	G0GCTG0GGA	AGAACA0AAA	CTG00CATCA	ACGCACTGAG	7600
CAACT0GTTG	CTA0G0CATC	ACAATCTGGT	GTATT0CA0C	ACTTCA0GCA	7650
GTCGTTG0CA	AAGGCAGAAG	AAAGTCACAT	TTGACAGACT	GCAAGTTCTG	7700
GACAG0CATT	A0CAGGACGT	GCTCAAGGAG	GTCAAAGCAG	CGGCGTCAAA	7750
AGTGAAGGCT	AAC TTGCTAT	00GTAGAGGA	AGCTTG0AGC	CTGACG000C	7800
CACATTCAGC	CAAATCCAAG	TTTGGCTATG	GGGCA0AAGA	CGTCCGTTGC	7850
CATG0CAGAA	AGG00GTAGC	CCACATCAAC	T00GTGTGGA	AAGACCTTCT	7900
GGAGACAGT	GTAA0ACCAA	TAGACACTAC	CATCATGG0C	AAGA0CGAGG	7950
TTTTCTG0GT	TCAG0CTGAG	AAGGGGGGTC	GTAA0CCAGC	T0GTCTCATC	8000
GTTT00000G	A0CTGGG0GT	G0G0GTTGTC	GAGAAGATGG	00CTGTACGA	8050
0GTGGTTAGC	AAGCT0000C	TGG00GTGAT	GGGAAGCT0C	TACGGATT0C	8100
AATACTCA0C	AGGACAG0GG	GTTGAATTCC	T0GTGCAAGC	GTGGAAGTCC	8150
AAGTAAG00C	0GATGGGGTT	CT0GTATGAT	A000GCTGTT	TTGACTCCAC	8200
AGTCACTGAG	AGCGACAT0C	GTACGGAGGA	GGCAATTTAC	CAATGTTGTG	8250
A0CTGGAC0C	CCAAG0000C	GTTGG0CATCA	AGT00CTCAC	TGAGAGGCTT	8300
TATGTTGGGG	G00CTCTTAC	CAATTCAAGG	GGGGAA0ACT	G0GGCTAC0G	8350
CAGGTG000C	G0GAG0GG0G	TACTGACAAC	TAGCTGTGGT	AACA00CTCA	8400
CTTGCTACAT	CAAGG0000G	GCAG0CTGTC	GAG00GCAGG	GCTCCAGGAC	8450
TGCA0CATGC	T0GTGTGTGG	0GAC0ACTTA	GTCGTTATCT	GTGAAAGTGC	8500
GGGGGT0CAG	GAGGACG0GG	0GAG0CTGAG	AG0CTTCACG	GAGGCTATGA	8550
0CAGGTACTC	0G0000000C	GGGGA0000C	CACA0CCAGA	ATACGACTTG	8600
GAGCTTATAA	CATCATGCTC	CTCCA0GTTG	TCAGT0G00C	ACGACGG0GC	8650
TGGAAAGAGG	GTCTACTACC	TTA000GTGA	00CTACAACC	000CT0G0GA	8700
GAG00G0GTG	GGAGACAGCA	AGACACACTC	CAGTCAATTC	CTGGCTAGGC	8750
AACATAATCA	TGTTTG000C	CACACTGTGG	G0GAGGATGA	TACTGATGAC	8800
0CATTTCCTT	AG0GT0CTCA	TAG0CAGGGA	TCAGCTTGAA	CAGGCTCTTA	8850
ACTGTGAGAT	CTACGGAG0C	TGCTACT0CA	TAGA0CCACT	GGATCTA0CT	8900
0CAATCATTC	AAAGACT0CA	TGG0CTCAGC	GCAATTTTAC	TCCACAGTTA	8950
CTCT0CAGGT	GAAATCAATA	GGGTGG000C	ATG0CTCAGA	AAACTTGGGG	9000

FIG. 1F

10	20	30	40	50	
1234567890	1234567890	1234567890	1234567890	1234567890	
TCCCGCCCTT	GCGAGCTTGG	AGACACCGGG	CCCGGAGCGT	CCGCGCTAGG	9050
CTTCTGTCCA	GAGGAGGCAG	GGCTGOCATA	TGTGGCAAGT	ACCTCTTCAA	9100
CTGGGCAGTA	AGAACAAAGC	TCAAACACAC	TCCAATAGCG	GCCGCTGGCC	9150
GGCTGGACTT	GTCCGGTTGG	TTCAACGGCTG	GCTACAGCGG	GGGAGACATT	9200
TATCACAGCG	TGTCATCATG	CCGGCCCCGC	TGGTCTCGGT	TTTGCCCTACT	9250
CCTGCTCCGT	GCAGGGGTAG	GCATCTACCT	CCGCCCAAC	CGATGAAGGT	9300
TGGGGTAAAC	ACTCCGGGCT	CTTAAGGCAT	TTCCGTGTTT	TTTTTTTTTT	9350
TTTTTTTTTT	TTTCTTTTTT	TTTTTCTTTT	CCTTTCCTTC	TTTTTTTCTT	9400
TTCTTTTTTC	CTTCTTTAAT	GGTGGCTCCA	TCCTAGCCCT	AGTCACGGCT	9450
AGCTGTGAAA	GGTCCGTGAG	CCGCATGACT	GCAGAGAGTG	CTGATACTGG	9500
CCTCTCTGCA	GATCATGT				9518

FIG. 1G

10	20	30	40	50	
1234567890	1234567890	1234567890	1234567890	1234567890	
MSINPKPQRK	TKRNINRRPQ	DVKFPGGGQI	VGGVYLLPRR	GPRLGVRATR	50
KTSERSQPRG	RROPIPKARR	PEGRTWAQPG	YPWPLYGNEG	CGWAGWLLSP	100
RGSRPSWGPT	DPRRRSRNLG	KVIDILITCGF	ADLMGYIPLV	GAPLGGAARA	150
LAHGVRVLED	GVNYATGNLP	GCSFSIFLLA	LLSCLITVPAS	AYQVRNSSGL	200
YHVINDCPNS	SIVYEADAI	LHTPGCVPCV	REGNASRCWV	AVIPTIVAIRD	250
GKLPITQLRR	HIDLLVGSAT	LCSALYVGDL	CGSVFLVGQL	FTFSPRRHWT	300
TQDQNCSTYP	GHTTGHMAW	DMMNWSPTA	ALVVAQLLRI	PQAIMDMIAG	350
AHWGVLAGIA	YFSMVGWNAK	VLVVLLLFAG	VDALQLININ	GSWHINSTAL	400
NONESLNTGW	LAGLFYQHKF	NSSGCPERLA	SCRRLTDFAQ	GWGPISYANG	450
SGLDERPYCW	HYPFRPGIV	PAKSVCGPVY	CFTSPFVVG	TTDRSGAPTY	500
SWGANDIDVF	VLNIRPPLG	NWFGCTWMNS	TGFTKVGAP	PCVIGGVGNN	550
TLLCPIDCFR	KHPEATYSRC	GSGPWITPRC	MVDYPYRLWH	YPCTINYTIF	600
KVRMYVGGVE	HRLEAACNWT	RGERCDLETR	DRSELSPLLL	STTQWQLPC	650
SFTTLPALST	GLIHLHQNTV	DVQYLYGVGS	SIASWAIKWE	YVLLFLLLA	700
DARVCSCLWM	MLLISQAEAA	LENLVIINAA	SLAGTHGLVS	FLVFFCFAWY	750
LKGRWPGAV	YALYGMWPLL	LLLLALPQRA	YALDTEVAAS	CGGVVLVGLM	800
ALITLSPYYKR	YLSWQMWLQ	YFLITRVEAQL	HWVWPLNVR	GGRDAVILLM	850
CVVHTILVFD	ITKLLLAIFG	PLWILQASLL	KVPYFVRVQG	LLRICALARK	900
IAGGHVQMA	ITKLGALTGT	YVYNHILTPLR	DWAHNGLRDL	AVAVEPVVFS	950
RMEIKLITWG	ADTAACGDII	NGLPVSARRG	QETLLGPADG	MVSKGWRLLA	1000
PITAYAQQTR	GLLGCIITSL	TGRDKNQVEG	EVQIVSTATQ	TFLATCINGV	1050
QWIVYHGAGT	RTIASPKGPV	IQMYTINVDQD	LVGWPAPOGS	RSLTPTCTGS	1100
SDLYLVIRHA	DVIPVRRRGD	SRGSLLSPRP	ISYLKGSSGG	PLTCPAGHAV	1150
GLFRAAVCTR	GVAKAVDFIP	VENLGTIMRS	PVFTDNSSPP	AVPQSFQVAH	1200
LHAPTGSCKS	TKVPAAYAAQ	GYKVLVLNPS	VAATLGFGAY	MSKAHVNDEN	1250
IRIGVRIITTT	GSPITYSTYG	KFLADGGCSG	GAYDIIICDE	CHSIDATSIL	1300
GIGIVLDQAE	TAGARLVWLA	TATPPGSVTV	SHNIEEVAL	STTGETIPFYG	1350
KAITPLEVIKG	GRHLIFCHSK	KKCELAACL	VALGINAVAY	YRGLDVSVIP	1400
TSGDVVVVST	DAIMIGFTGD	FDSVIDCNTC	VIQIVDFSLD	PFTTIEITIL	1450
PQDAVSRIQR	RGRIGRGKPG	TYRFVAPGER	PSGMFDSSVL	CECYDAGCAW	1500
YELTPAETIV	RLRAYMNTFG	LPVQQDHLEF	WEGVFTGLTH	IDAHLFSQIK	1550
QSGENFPYLV	AYQATVCARA	QAPPPSWDQM	WKCLIRLKPT	LHGPTPLLYR	1600
LGAVQNEVIL	THPTIKYIMT	CMSADLEVVT	STWLVGGVL	AALAAAYCLST	1650
GCVWIVGRIV	LSGKPAIIPD	REVLVQEFDE	MEECSQHLPY	IBQGMMLAEQ	1700
FKQKALGLIQ	TASRHAEVIT	PAVQINWQKL	EVFWAKHMAN	FISGIQYLAG	1750
LSTLPGNPAI	ASLMAFTAAV	TSPLITGQTL	LFNILLGGWA	AQLAAPGAAT	1800
AFVGAAGLAGA	AIGSVGLGKV	LVDILAGYGA	GVAGALVAFK	IMSGEVPSTE	1850
DLVNLLPAIL	SFGALWGVV	CAAILRRHVG	PGEQAVQWMN	RLIAFASRGN	1900

FIG. 1H

10	20	30	40	50	
1234567890	1234567890	1234567890	1234567890	1234567890	
HVSPTHYVPE	SDAAARVTAI	LSSLIVTQLL	RRLHQWISSE	CTTFCSGSWL	1950
RDIWDWICEV	LSDFKIWLKA	KLMPQLPGIP	FVSCQRGYRG	VWRGDGIMHT	2000
RCHOGAETTG	HVKNGIMRIV	GPRICRMMS	GIFPINAYTT	GPCTPLPAEN	2050
YKFALWRVSA	EEYVEIRRVG	DFHVSGMIT	DNLCPCQIIP	SPEFFTELDG	2100
VRLHRFAPPC	KPLLREEVSF	RVGLHEYPVG	SQLPCEPEPD	VAVLTSMLTD	2150
PSHTTAEAG	RRLARGSPPS	MASSASQLS	APSLKATCTA	NHDSFDAELI	2200
EANLLWRQEM	GGNIRVESE	NKWILDSFD	PLVAEEDERE	VSVPAEILRK	2250
SRRFARALPV	WARDYNPPL	VEIWKKPDYE	PPVHGCPLP	PPRSPPVPPP	2300
RKKRTVVLTE	STLSTALAE	ATKSFGSSST	SGTIGINTTT	SSEPAPSGCP	2350
PDSDESYSYSS	MPPLEGETPD	PDLSDGSWST	VSSGADTEIV	VCCSMSYSWT	2400
GALVTPCAEE	EQKLPINALS	NSLLRHHNLV	YSTTSRSACQ	RQKKVTFDRL	2450
QVLDSHYQDV	LKEVKAAASK	VKANLLSVEE	ACSLTPPHSA	KSKFGYGAKD	2500
VRCHARKAVA	HINSVWKDLL	EDSVTPIDTT	IMAKNEVFCV	QPEKGGRKPA	2550
RLIVFPDLGV	RVCEKMALYD	VSKLPLAVM	GSSYGFQYSP	GQRVEFLVQA	2600
WKSCKTPMGF	SYDIRCFDST	VTESDIRTEE	AIYQCCDLDP	QARVAIKSLT	2650
ERLYVGGPLT	NSRGENCGYR	RCRASGVLT	SGNITLCYI	KARAACRAAG	2700
LQDCIMLVCG	DDLWICESA	GVQEDAASLR	AFTEAMIRYS	APPGDPPQPE	2750
YDLELITSCS	SNVSVAHGDA	GKRVYYLIRD	PTTPLARAAW	ETARHTPVNS	2800
WLGNIIMFAP	TLWARMILMT	HFFSVLIARD	QLEQALNCEI	YGACYSIEPL	2850
DLPTTIQRLH	GLSAFSLHSY	SPGETNRVAA	CLRKLGVPPL	RAWRHRARSV	2900
RARLLSRGGR	AAICGKYLEN	WAVRIKLIKLT	PIAAAGRLLD	SGWFTAGYSG	2950
GDIYHSVSHA	RPRWFWFCLL	LLAAGVGIYL	LPNR		2984

FIG. 2

In Vitro Transcription Translation

Lane	1	3	4	5	6	8
Canine Membranes	-	+	++	+++	++++	++




FIG. 3

Chimpanzee 1590: Transfection with RNA Transcripts of
Hepatitis C Virus HVR1 Deletion Mutant





Figure 4

Infection of Chimpanzee 96A008 with Δ HVR1



SEQUENCE LISTING

<110> FORNS, XAVIER
 BUKH, JENS
 EMERSON, SUZANNE U.
 PURCELL, ROBERT H.

<120> NUCLEIC ACID MOLECULES ENCODING HEPATITIS C VIRUS WHICH
 LACK HYPERVARIABLE REGION ONE OF E2 AND USES THEREOF

<130> 2026-431PC

<140> To Be Assigned

<141> 2000-09-22

<150> 60/155,823

<151> 1999-09-23

<160> 12

<170> PatentIn Ver. 2.1

<210> 1

<211> 9518

<212> DNA

<213> Hepatitis C virus

<400> 1

```

gccagccccc tgatgggggc gacactccac catgaatcac tcccctgtga ggaactactg 60
tcttcacgca gaaagcgtct agccatggcg ttagtatgag tgtcgtgcag cctccaggac 120
ccccctccc gggagagcca tagtgggtctg cggaaccggt gagtacaccg gaattgccag 180
gacgaccggg tcctttcttg gataaacccg ctcaatgcct ggagatttgg gcgtgcccc 240
gcaagactgc tagccgagta gtgttgggtc gcgaaaggcc ttgtggtact gcctgatagg 300
gtgcttgcca gtgccccggg aggtctcgta gaccgtgcac catgagcacg aatcctaaac 360
ctcaaagaaa aaccaaacgt aacaccaacc gtgcccaca ggacgtcaag ttcccgggtg 420
gcggtcagat cgttgggtgga gtttacttgt tgccgcgcag gggccctaga ttgggtgtgc 480
gcgcgacgag gaagacttcc gagcggtcgc aacctcgagg tagacgtcag cctatcccca 540
aggcacgtcg gcccaggggc aggacctggg ctgagcccgg gtacccttgg cccctctatg 600
gcaatgaggg ttgcgggttg gcgggatggc tcctgtctcc cgtggtctct cggcctagct 660
ggggccccc agacccccgg cgtaggtcgc gcaatttggg taaggtcac gataccctta 720
cgtgcggctt cgccgacctc atggggtaca taccgtctgt cggcgccctt cttggaggcg 780
ctgccagggc cctggcgcat ggcgtccggg ttctggaaga cggcgtgaac tatgcaacag 840
ggaaccttcc tggttgctct ttctctatct tccttctggc cctgctctct tgccctgactg 900
tgcccgcctc agcctaccaa gtgcgcaatt cctcggggct ttaccatgtc accaatgatt 960
gccctaactc gagtattgtg tacgaggcgg ccgatgccat cctgcacact ccgggggtgtg 1020
tcccttgcgt tcgcgagggg aacgcctcga ggtgttgggt ggcggtgacc cccacggtgg 1080
ccaccaggga cggcaaacct ccacaacgc agcttcgacg tcatatcgat ctgcttctgc 1140
ggagcgccac cctctgctcg gccctctacg tgggggacct gtgcgggtct gtctttcttg 1200

```

```

ttggtcaact gtttaccttc tctcccaggc gccactggac gacgcaagac tgcaattggt 1260
ctatctatcc cggccatata acgggtcatc gcatggcatg ggatatgatg atgaactggg 1320
cccctacggc agcgttggtg gtagctcagc tgctccggat cccacaagcc atcatggaca 1380
tgatcgctgg tgctcactgg ggagtcctgg cgggcatagc gtatttctcc atggtgggga 1440
actgggcgaa ggtcctggta gtgctgctgc tatttgccgg cgtcgacgcg atccaactga 1500
tcaacaccaa cggcagttgg cacatcaata gcacggcctt gaattgcaat gaaagcctta 1560
acaccggctg gttagcaggg ctcttctatc aacacaaatt caactcttca ggctgtcctg 1620
agaggttggc cagctgccga cgcttaccg attttgccca gggctggggg cctatcagtt 1680
atgccaacgg aagcggcctc gacgaacgcc cctactgctg gcactaccct ccaagacctt 1740
gtggcattgt gcccgcaaag agcgtgtgtg gcccgggata ttgcttctc cccagccccg 1800
tggtgggtgg aacgaccgac aggtcggggc cgcctaccta cagctggggg gcaaatgata 1860
cggatgtctt cgtccttaac aacaccaggc caccgctggg caattgggtt ggttgtaacct 1920
ggatgaactc aactggattc accaaagtgt gcggagcgcc cccttggtgc atcggagggg 1980
tgggcaacaa cacttgctc tgccccactg attgcttccg caaacatccg gaagccacat 2040
actctcggtg cggctccggt ccctggatta caccaggtg catggtcgac taccctgata 2100
ggctttggca ctatccttgt accatcaatt acaccatatt caaagtcagg atgtacgtgg 2160
gaaggggtcga gcacaggctg gaagcggcct gcaactggac gcggggcgaa cgctgtgatc 2220
tggaagacag ggacaggctc gagctcagcc cgttgctgct gtccaccaca cagtggcagg 2280
tccttccgtg ttcttccacg accctgccag ccttgctcac cggectcatc cacctccacc 2340
agaacattgt ggacgtgcag tacttgtagc gggtagggtc aagcatcgcg tcctgggcca 2400
ttaagtggga gtacgtcggt ctctgttcc ttctgcttgc agacgcgcgc gtctgtcctc 2460
gcttggtgat gatgtactc atatcccaag cggaggcggc tttggagaac ctcgtaatac 2520
tcaatgcagc atccctggcc gggacgcacg gtcttggtgc ctctctcgtg ttcttctgct 2580
ttgcgtggta tctgaagggt aggtgggtgc ccggagcggc ctacgccctc tacgggatgt 2640
ggcctctcct cctgtcctc ctggcggtgc ctacgcggc atacgcactg gacacggagg 2700
tgcccgctgc gtgtggcggc gttgttcttg tggggttaat ggcgtgact ctgtcgccat 2760
attacaagcg ctatcagcag tggtagcatg ggtggcttca gtatttctg accagagtag 2820
aagcgcaact gcacgtgtgg gttccccccc tcaacgtccg gggggggcgc gatgccgtca 2880
tcttactcat gtgtgtagta cacccgaccc tggattttga catcaccaaa ctactcctgg 2940
ccatcttcgg acccctttgg attcttcaag ccagtttgct taaagtcccc tacttctgtc 3000
gcgttcaagg cttctcctcg atctgcgcgc tagcgcggaa gatagccgga ggtcattacg 3060
tgcaaatggc catcatcaag ttaggggcgc ttactggcac ctatgtgat aaccatctca 3120
cccctcttcg agactgggag cacaacggcc tgcgagatct ggccgtggct gtggaaccag 3180
tcgtcttctc ccgaatggag accaagctca tcacgtgggg ggcagatacc gccgcgtgct 3240
gtgacatcat caacggcctg cccgtctctg cccgtagggg ccaggagata ctgcttgggc 3300
cagccgacgg aatgggtctc aaggggtgga ggttgctggc gccatcacg gcgtacgccc 3360
agcagacgag aggcctccta ggggtgtataa tcaccagcct gactggccgg gacaaaaacc 3420
aagtggaggg tgaggtccag atcgtgtcaa ctgctaccca aaccttctg gcaacgtgca 3480
tcaatggggg atgctggact gtctaccacg gggccggaac gaggaccatc gcatcaccca 3540
agggctcctg catccagatg tataccaatg tggaccaaga ccttggtggc tggcccgctc 3600
ctcaagggtc ccgtcattg acaccctgta cctgcggctc ctcgacctt tacctgggtc 3660
cgaggcacgc cgatgtcatt cccgtgcgcc ggcgaggtga tagcaggggt agcctgcttt 3720
cgccccggcc catttctac ttgaaaggct cctcgggggg tccgctgttg tgcccccgcg 3780
gacacgccgt gggcctattc agggccgcgg tgtgcacccg tggagtggct aaagcgggtg 3840
actttatccc tgtggagaac ctagggacaa ccatgagatc cccggtgttc acggacaact 3900
cctctccacc agcagtgcac cagagcttcc aggtggccca cctgcatgct cccaccggca 3960
gcggtaagag caccaaggct ccggctgcgt acgcagccca gggctacaag gtgttggtgc 4020
tcaaccctc tggtgctgca acgctgggct ttggtgctta catgtccaag gcccatgggg 4080

```

```

ttgatacctaa tatcaggacc ggggtgagaa caattaccac tggcagcccc atcacgtact 4140
ccacctacgg caagttcctt gccgacggcg ggtgctcagg aggtgcttat gacataataa 4200
tttgtgacga gtgccactcc acggatgcca catccatctt gggcatcggc actgtccttg 4260
accaagcaga gactgcgggg gcgagactgg ttgtgctcgc cactgctacc cctccgggct 4320
ccgtcactgt gtcccatcct aacatcgagg aggttgcctt gtccaccacc ggagagatcc 4380
ccttttacgg caaggctatc cccctcgagg tgatcaaggg ggggaagacat ctcatcttct 4440
gccactcaaa gaagaagtgc gacgagctcg ccgcgaagct ggtcgcattg ggcatcaatg 4500
ccgtggccta ctaccgcggg cttgacgtgt ctgtcatccc gaccagcggc gatgttgctg 4560
tcgtgtcgac cgatgctctc atgactggct ttaccggcga cttcgactct gtgatagact 4620
gcaaacctg tgctactcag acagtgcatt tcagccttga ccctacctt accattgaga 4680
caaccacgct ccccgaggat gctgtctcca ggactcaacg ccggggcagg actggcaggg 4740
ggaagccagg catctataga tttgtggcac cgggggagcg cccctccggc atgttcgact 4800
cgtccgtcct ctgtgagtgc tatgacgcgg gctgtgcttg gtatgagctc acgcccgcg 4860
agactacagt taggtacga gcgtacatga acaccccggg gcttcccgtg tggcaggacc 4920
atcttgaatt ttgggagggc gtctttacgg gcctcactca tatagatgcc cactttttat 4980
cccagacaaa gcagagtggg gagaactttc cttacctggt agcgtaccaa gccaccgtgt 5040
gcgctagggc tcaagccctt ccccatcggt gggaccagat gtggaagtgt ttgatccgcc 5100
ttaaacccac cctccatggg ccaacacccc tgctatacag actgggcgct gttcagaatg 5160
aagtcaccct gacgcaccca atcaccaa atcatcatgac atgcatgtcg gccgacctgg 5220
aggctgtcac gagcacctgg gtgctcggtt gcggcgctct ggctgctctg gccgcgtatt 5280
gcctgtcaac aggtgcgctg gtcatagtgg gcaggatcgt cttgtccggg aagccggcaa 5340
ttatacctga caggagggtt ctctaccagg agttcgatga gatggaagag tgctctcagc 5400
acttacgtga catcgagcaa gggatgatgc tcgctgagca gttcaagcag aaggccctcg 5460
gcctcctgca gaccgcgtcc cgcatgcag aggttatcac ccctgctgtc cagaccaact 5520
ggcagaaaact cgaggctctt tgggcgaagc acatgtggaa tttcatcagt gggatacaat 5580
acttggcggg cctgtcaacg ctgcctggta accccgccat tgcttcattg atggctttta 5640
cagctgccgt caccagccca ctaaccactg gccaaaccct cctcttcaac atattggggg 5700
ggtgggtggc tgcccagctc gccgcccccg gtgcccgtac tgcccttggt ggtgctggcc 5760
tagctggcgc cgccatcggc agcgttggac tggggaaggt cctcgtggac attcttgacg 5820
ggtatggcgc gggcgtggcg ggagctcttg tagcattcaa gatcatgagc ggtgaggtcc 5880
cctccacgga ggacctggtc aatctgctgc ccgccatcct ctgcctgga gcccttgtag 5940
tcggtgtggt ctgctgagca atactgcgcc ggcacgttgg cccgggcgag ggggcagtcg 6000
aatggatgaa ccggctaata gccttcgcct cccgggggaa ccatgtttcc cccacgcact 6060
acgtgccgga gagcgatgca gccgcccgcg tcaactgcat actcagcagc ctactgttaa 6120
cccagctcct gaggcgactg catcagtggg taagctcgga gtgtaccact ccatgctccg 6180
gttcctggct aaggacatc tgggactgga tatgagaggt gctgagcgac tttaaacct 6240
ggctgaaagc caagctcatg ccacaactgc ctgggattcc ctttgtgtcc tgccagcgcg 6300
ggtatagggg ggtctggcga ggagacggca ttatgcacac tcgctgccac tgtggagctg 6360
agatcactgg acatgtcaaa aacgggacga tgaggatcgt cggctcctagg acctgcagga 6420
acatgtggag tgggacgttc cccattaacg cctacaccac gggccctgt actccccctc 6480
ctgcgccgaa ctataagttc gcgctgtgga ggggtgtctg agaggaatac gtggagataa 6540
ggcgggtggg ggacttccac tacgtatcgg gtatgactac tgacaatctt aaatgcccgt 6600
gccagatccc atcgcccgaa tttttcacag aattggacgg ggtgcgccta cacaggtttg 6660
cgcccccttg caagcccttg ctgcgggagg aggtatcatt cagagtagga ctccacgagt 6720
accgggtggg gtcgcaatta ccttgcgagc ccgaaccgga cgtagccgtg ttgacgtcca 6780
tgctcactga tccctcccat ataacagcag agggggccgg gagaaggttg gcgagaggg 6840
cacccttc tatggccagc tcctcggtta gccagctgtc cgctccatct ctcaaggcaa 6900
cttgaccgc caaccatgac tcccctgacg ccgagctcat agaggctaac ctctgtgga 6960

```

```

ggcaggagat gggcggaac atcaccaggg ttgagtcaga gaacaaagtgt gtgattctgtg 7020
actccttcga tccgcttggt gcagaggagg atgagcggga ggtctccgta cctgcagaaa 7080
ttctgcggaa gtctcggaga ttcgcccggg ccctgcccgt ctgggcgcgg ccggactaca 7140
accccccgct agtagagacg tggaaaaagc ctgactacga accacctgtg gtccatggct 7200
gcccgtacc acctccacgg tcccctcctg tgcctccgcc tcggaaaaag cgtacgggtg 7260
tcctcaccga atcaacccta tctactgect tggccgagct tgccaccaa agttttggca 7320
gctcctcaac ttccggcatt acgggcgaca atacgacaac atcctctgag cccgccccct 7380
ctggctgccc cccgactcc gacgttgagt cctattcttc catgccccct ctggaggggg 7440
agcctgggga tccggatctc agcgacgggt catggtcgac ggtcagtagt ggggcggaca 7500
cgggaagatgt cgtgtgctgc tcaatgtctt attcctggac aggcgcactc gtcaccccg 7560
gcgctgcgga agaacaacaa ctgcccata acgcactgag caactcgttg ctacgccatc 7620
acaatctggt gtattccacc acttcacgca gtgcttgcca aaggcagaag aaagtcacat 7680
ttgacagact gcaagttctg gacagccatt accaggacgt gctcaaggag gtcaaagcag 7740
cggcgctcaaa agtgaaggct aacttgctat ccgtagagga agcttgacgc ctgacgcccc 7800
cacattcagc caaatccaag tttggctatg gggcaaaaaga cgtccgttgc catgccagaa 7860
aggccgtagc ccacatcaac tccgtgtgga aagaccttct ggaagacagt gtaacaccaa 7920
tagacactac catcatggcc aagaacgagg ttttctgcgt tcagcctgag aaggggggtc 7980
gtaagccagc tcgtctcacc gtgttccccg acctgggggt gcgctgtgca gagaagatgg 8040
ccctgtacga cgtggttagc aagctcccc tggccgtgat gggagactcc tacggattcc 8100
aatactcacc aggacagcgg gttgaattcc tcgtgcaagc gtggaagtcc aagaagaccc 8160
cgatgggggt ctcgtatgat acccgctgtt ttgactccac agtcaactgag agcgacatcc 8220
gtacggagga ggcaatttac caatgttgtg acctggacct ccaagccgc gtggccatca 8280
agtccctcac tgagaggctt tatgttgggg gccctcttac caattcaagg ggggaaaaact 8340
gcggctaccg caggtgccgc gcgagcggcg tactgacaac tagctgtggt aacaccctca 8400
cttgtctacat caaggcccg gcagcctgtc gagccgcagg gctccaggac tgcaccatgc 8460
tcgtgtgtgg cgacgactta gtcgttatct gtgaaagtgc ggggggtccag gaggacgcgg 8520
cgagcctgag agccttcacg gaggctatga ccaggctact cggccccccc ggggaccccc 8580
cacaaccaga atacgacttg gagcttataa catcatgctc ctccaacgtg tcagtgcgcc 8640
acgagggcgc tggaaagagg gtctactacc ttaccctgta ccctacaacc cccctcgcca 8700
gagccgcgtg ggagacagca agacacactc cagtcaattc ctggctaggc aacataatca 8760
tgtttgcccc cacactgtgg gcgaggatga tactgatgac ccatttcttt agcgtcctca 8820
tagccaggga tcagcttgaa caggctctta actgtgagat ctacggagcc tgctactcca 8880
tagaaccact ggatctacct ccaatcattc aaagactcca tggcctcagc gcattttcac 8940
tccacagtta ctctccaggt gaaatcaata ggggtggcgc atgcctcaga aaacttgggg 9000
tcccgcctt gcgagcttg agacacggg cccggagcgt ccgcgctagg cttctgtcca 9060
gaggaggcag ggctgccata tgtggcaagt acctcttcaa ctgggcagta agaacaaagc 9120
tcaaactcac tccaatagcg gccgctggcc ggctggactt gtccggttgg ttcacggctg 9180
gctacagcgg gggagacatt tatcacagcg tgtctcatgc ccggccccgc tggttctggt 9240
tttgctact cctgctcgct gcaggggtag gcatctacct cctcccaac cgatgaagg 9300
tggggtaaac actccggcct cttaagccat ttctgtttt ttttttttt ttttttttt 9360
ttttctttt tttttcttt ctttctctt ttttttctt ttctttttt cttctttaat 9420
ggtggctcca tcttagccct agtcacggct agctgtgaaa ggtccgtgag ccgcatgact 9480
gcagagagtg ctgatactgg cctctctgca gatcatgt 9518

```

<210> 2

<211> 2984

<212> PRT

<213> Hepatitis C virus

<400> 2

```

Met Ser Thr Asn Pro Lys Pro Gln Arg Lys Thr Lys Arg Asn Thr Asn
  1              5              10              15

Arg Arg Pro Gln Asp Val Lys Phe Pro Gly Gly Gly Gln Ile Val Gly
      20              25              30

Gly Val Tyr Leu Leu Pro Arg Arg Gly Pro Arg Leu Gly Val Arg Ala
      35              40              45

Thr Arg Lys Thr Ser Glu Arg Ser Gln Pro Arg Gly Arg Arg Gln Pro
      50              55              60

Ile Pro Lys Ala Arg Arg Pro Glu Gly Arg Thr Trp Ala Gln Pro Gly
      65              70              75              80

Tyr Pro Trp Pro Leu Tyr Gly Asn Glu Gly Cys Gly Trp Ala Gly Trp
      85              90              95

Leu Leu Ser Pro Arg Gly Ser Arg Pro Ser Trp Gly Pro Thr Asp Pro
      100             105             110

Arg Arg Arg Ser Arg Asn Leu Gly Lys Val Ile Asp Thr Leu Thr Cys
      115             120             125

Gly Phe Ala Asp Leu Met Gly Tyr Ile Pro Leu Val Gly Ala Pro Leu
      130             135             140

Gly Gly Ala Ala Arg Ala Leu Ala His Gly Val Arg Val Leu Glu Asp
      145             150             155             160

Gly Val Asn Tyr Ala Thr Gly Asn Leu Pro Gly Cys Ser Phe Ser Ile
      165             170             175

Phe Leu Leu Ala Leu Leu Ser Cys Leu Thr Val Pro Ala Ser Ala Tyr
      180             185             190

Gln Val Arg Asn Ser Ser Gly Leu Tyr His Val Thr Asn Asp Cys Pro
      195             200             205

Asn Ser Ser Ile Val Tyr Glu Ala Ala Asp Ala Ile Leu His Thr Pro
      210             215             220

Gly Cys Val Pro Cys Val Arg Glu Gly Asn Ala Ser Arg Cys Trp Val
      225             230             235             240

```

Ala	Val	Thr	Pro	Thr	Val	Ala	Thr	Arg	Asp	Gly	Lys	Leu	Pro	Thr	Thr				
				245						250						255			
Gln	Leu	Arg	Arg	His	Ile	Asp	Leu	Leu	Val	Gly	Ser	Ala	Thr	Leu	Cys				
				260						265						270			
Ser	Ala	Leu	Tyr	Val	Gly	Asp	Leu	Cys	Gly	Ser	Val	Phe	Leu	Val	Gly				
				275						280						285			
Gln	Leu	Phe	Thr	Phe	Ser	Pro	Arg	Arg	His	Trp	Thr	Thr	Gln	Asp	Cys				
				290						295						300			
Asn	Cys	Ser	Ile	Tyr	Pro	Gly	His	Ile	Thr	Gly	His	Arg	Met	Ala	Trp				
305						310						315						320	
Asp	Met	Met	Met	Asn	Trp	Ser	Pro	Thr	Ala	Ala	Leu	Val	Val	Ala	Gln				
				325						330						335			
Leu	Leu	Arg	Ile	Pro	Gln	Ala	Ile	Met	Asp	Met	Ile	Ala	Gly	Ala	His				
				340						345						350			
Trp	Gly	Val	Leu	Ala	Gly	Ile	Ala	Tyr	Phe	Ser	Met	Val	Gly	Asn	Trp				
				355						360						365			
Ala	Lys	Val	Leu	Val	Val	Leu	Leu	Leu	Phe	Ala	Gly	Val	Asp	Ala	Ile				
				370						375						380			
Gln	Leu	Ile	Asn	Thr	Asn	Gly	Ser	Trp	His	Ile	Asn	Ser	Thr	Ala	Leu				
385						390						395						400	
Asn	Cys	Asn	Glu	Ser	Leu	Asn	Thr	Gly	Trp	Leu	Ala	Gly	Leu	Phe	Tyr				
				405						410						415			
Gln	His	Lys	Phe	Asn	Ser	Ser	Gly	Cys	Pro	Glu	Arg	Leu	Ala	Ser	Cys				
				420						425						430			
Arg	Arg	Leu	Thr	Asp	Phe	Ala	Gln	Gly	Trp	Gly	Pro	Ile	Ser	Tyr	Ala				
				435						440						445			
Asn	Gly	Ser	Gly	Leu	Asp	Glu	Arg	Pro	Tyr	Cys	Trp	His	Tyr	Pro	Pro				
				450						455						460			
Arg	Pro	Cys	Gly	Ile	Val	Pro	Ala	Lys	Ser	Val	Cys	Gly	Pro	Val	Tyr				
465						470						475						480	
Cys	Phe	Thr	Pro	Ser	Pro	Val	Val	Val	Gly	Thr	Thr	Asp	Arg	Ser	Gly				
				485						490						495			

Ala Pro Thr Tyr Ser Trp Gly Ala Asn Asp Thr Asp Val Phe Val Leu
 500 505 510

Asn Asn Thr Arg Pro Pro Leu Gly Asn Trp Phe Gly Cys Thr Trp Met
 515 520 525

Asn Ser Thr Gly Phe Thr Lys Val Cys Gly Ala Pro Pro Cys Val Ile
 530 535 540

Gly Gly Val Gly Asn Asn Thr Leu Leu Cys Pro Thr Asp Cys Phe Arg
 545 550 555 560

Lys His Pro Glu Ala Thr Tyr Ser Arg Cys Gly Ser Gly Pro Trp Ile
 565 570 575

Thr Pro Arg Cys Met Val Asp Tyr Pro Tyr Arg Leu Trp His Tyr Pro
 580 585 590

Cys Thr Ile Asn Tyr Thr Ile Phe Lys Val Arg Met Tyr Val Gly Gly
 595 600 605

Val Glu His Arg Leu Glu Ala Ala Cys Asn Trp Thr Arg Gly Glu Arg
 610 615 620

Cys Asp Leu Glu Asp Arg Asp Arg Ser Glu Leu Ser Pro Leu Leu Leu
 625 630 635 640

Ser Thr Thr Gln Trp Gln Val Leu Pro Cys Ser Phe Thr Thr Leu Pro
 645 650 655

Ala Leu Ser Thr Gly Leu Ile His Leu His Gln Asn Ile Val Asp Val
 660 665 670

Gln Tyr Leu Tyr Gly Val Gly Ser Ser Ile Ala Ser Trp Ala Ile Lys
 675 680 685

Trp Glu Tyr Val Val Leu Leu Phe Leu Leu Leu Ala Asp Ala Arg Val
 690 695 700

Cys Ser Cys Leu Trp Met Met Leu Leu Ile Ser Gln Ala Glu Ala Ala
 705 710 715 720

Leu Glu Asn Leu Val Ile Leu Asn Ala Ala Ser Leu Ala Gly Thr His
 725 730 735

Gly Leu Val Ser Phe Leu Val Phe Phe Cys Phe Ala Trp Tyr Leu Lys
 740 745 750

Gly Arg Trp Val Pro Gly Ala Val Tyr Ala Leu Tyr Gly Met Trp Pro
 755 760 765
 Leu Leu Leu Leu Leu Leu Ala Leu Pro Gln Arg Ala Tyr Ala Leu Asp
 770 775 780
 Thr Glu Val Ala Ala Ser Cys Gly Gly Val Val Leu Val Gly Leu Met
 785 790 795 800
 Ala Leu Thr Leu Ser Pro Tyr Tyr Lys Arg Tyr Ile Ser Trp Cys Met
 805 810 815
 Trp Trp Leu Gln Tyr Phe Leu Thr Arg Val Glu Ala Gln Leu His Val
 820 825 830
 Trp Val Pro Pro Leu Asn Val Arg Gly Gly Arg Asp Ala Val Ile Leu
 835 840 845
 Leu Met Cys Val Val His Pro Thr Leu Val Phe Asp Ile Thr Lys Leu
 850 855 860
 Leu Leu Ala Ile Phe Gly Pro Leu Trp Ile Leu Gln Ala Ser Leu Leu
 865 870 875 880
 Lys Val Pro Tyr Phe Val Arg Val Gln Gly Leu Leu Arg Ile Cys Ala
 885 890 895
 Leu Ala Arg Lys Ile Ala Gly Gly His Tyr Val Gln Met Ala Ile Ile
 900 905 910
 Lys Leu Gly Ala Leu Thr Gly Thr Tyr Val Tyr Asn His Leu Thr Pro
 915 920 925
 Leu Arg Asp Trp Ala His Asn Gly Leu Arg Asp Leu Ala Val Ala Val
 930 935 940
 Glu Pro Val Val Phe Ser Arg Met Glu Thr Lys Leu Ile Thr Trp Gly
 945 950 955 960
 Ala Asp Thr Ala Ala Cys Gly Asp Ile Ile Asn Gly Leu Pro Val Ser
 965 970 975
 Ala Arg Arg Gly Gln Glu Ile Leu Leu Gly Pro Ala Asp Gly Met Val
 980 985 990
 Ser Lys Gly Trp Arg Leu Leu Ala Pro Ile Thr Ala Tyr Ala Gln Gln
 995 1000 1005

Thr Arg Gly Leu Leu Gly Cys Ile Ile Thr Ser Leu Thr Gly Arg Asp
 1010 1015 1020
 Lys Asn Gln Val Glu Gly Glu Val Gln Ile Val Ser Thr Ala Thr Gln
 1025 1030 1035 1040
 Thr Phe Leu Ala Thr Cys Ile Asn Gly Val Cys Trp Thr Val Tyr His
 1045 1050 1055
 Gly Ala Gly Thr Arg Thr Ile Ala Ser Pro Lys Gly Pro Val Ile Gln
 1060 1065 1070
 Met Tyr Thr Asn Val Asp Gln Asp Leu Val Gly Trp Pro Ala Pro Gln
 1075 1080 1085
 Gly Ser Arg Ser Leu Thr Pro Cys Thr Cys Gly Ser Ser Asp Leu Tyr
 1090 1095 1100
 Leu Val Thr Arg His Ala Asp Val Ile Pro Val Arg Arg Arg Gly Asp
 1105 1110 1115 1120
 Ser Arg Gly Ser Leu Leu Ser Pro Arg Pro Ile Ser Tyr Leu Lys Gly
 1125 1130 1135
 Ser Ser Gly Gly Pro Leu Leu Cys Pro Ala Gly His Ala Val Gly Leu
 1140 1145 1150
 Phe Arg Ala Ala Val Cys Thr Arg Gly Val Ala Lys Ala Val Asp Phe
 1155 1160 1165
 Ile Pro Val Glu Asn Leu Gly Thr Thr Met Arg Ser Pro Val Phe Thr
 1170 1175 1180
 Asp Asn Ser Ser Pro Pro Ala Val Pro Gln Ser Phe Gln Val Ala His
 1185 1190 1195 1200
 Leu His Ala Pro Thr Gly Ser Gly Lys Ser Thr Lys Val Pro Ala Ala
 1205 1210 1215
 Tyr Ala Ala Gln Gly Tyr Lys Val Leu Val Leu Asn Pro Ser Val Ala
 1220 1225 1230
 Ala Thr Leu Gly Phe Gly Ala Tyr Met Ser Lys Ala His Gly Val Asp
 1235 1240 1245
 Pro Asn Ile Arg Thr Gly Val Arg Thr Ile Thr Thr Gly Ser Pro Ile
 1250 1255 1260

Thr Tyr Ser Thr Tyr Gly Lys Phe Leu Ala Asp Gly Gly Cys Ser Gly
 1265 1270 1275 1280
 Gly Ala Tyr Asp Ile Ile Ile Cys Asp Glu Cys His Ser Thr Asp Ala
 1285 1290 1295
 Thr Ser Ile Leu Gly Ile Gly Thr Val Leu Asp Gln Ala Glu Thr Ala
 1300 1305 1310
 Gly Ala Arg Leu Val Val Leu Ala Thr Ala Thr Pro Pro Gly Ser Val
 1315 1320 1325
 Thr Val Ser His Pro Asn Ile Glu Glu Val Ala Leu Ser Thr Thr Gly
 1330 1335 1340
 Glu Ile Pro Phe Tyr Gly Lys Ala Ile Pro Leu Glu Val Ile Lys Gly
 1345 1350 1355 1360
 Gly Arg His Leu Ile Phe Cys His Ser Lys Lys Lys Cys Asp Glu Leu
 1365 1370 1375
 Ala Ala Lys Leu Val Ala Leu Gly Ile Asn Ala Val Ala Tyr Tyr Arg
 1380 1385 1390
 Gly Leu Asp Val Ser Val Ile Pro Thr Ser Gly Asp Val Val Val Val
 1395 1400 1405
 Ser Thr Asp Ala Leu Met Thr Gly Phe Thr Gly Asp Phe Asp Ser Val
 1410 1415 1420
 Ile Asp Cys Asn Thr Cys Val Thr Gln Thr Val Asp Phe Ser Leu Asp
 1425 1430 1435 1440
 Pro Thr Phe Thr Ile Glu Thr Thr Thr Leu Pro Gln Asp Ala Val Ser
 1445 1450 1455
 Arg Thr Gln Arg Arg Gly Arg Thr Gly Arg Gly Lys Pro Gly Ile Tyr
 1460 1465 1470
 Arg Phe Val Ala Pro Gly Glu Arg Pro Ser Gly Met Phe Asp Ser Ser
 1475 1480 1485
 Val Leu Cys Glu Cys Tyr Asp Ala Gly Cys Ala Trp Tyr Glu Leu Thr
 1490 1495 1500
 Pro Ala Glu Thr Thr Val Arg Leu Arg Ala Tyr Met Asn Thr Pro Gly
 1505 1510 1515 1520

Leu Pro Val Cys Gln Asp His Leu Glu Phe Trp Glu Gly Val Phe Thr
 1525 1530 1535
 Gly Leu Thr His Ile Asp Ala His Phe Leu Ser Gln Thr Lys Gln Ser
 1540 1545 1550
 Gly Glu Asn Phe Pro Tyr Leu Val Ala Tyr Gln Ala Thr Val Cys Ala
 1555 1560 1565
 Arg Ala Gln Ala Pro Pro Pro Ser Trp Asp Gln Met Trp Lys Cys Leu
 1570 1575 1580
 Ile Arg Leu Lys Pro Thr Leu His Gly Pro Thr Pro Leu Leu Tyr Arg
 1585 1590 1595 1600
 Leu Gly Ala Val Gln Asn Glu Val Thr Leu Thr His Pro Ile Thr Lys
 1605 1610 1615
 Tyr Ile Met Thr Cys Met Ser Ala Asp Leu Glu Val Val Thr Ser Thr
 1620 1625 1630
 Trp Val Leu Val Gly Gly Val Leu Ala Ala Leu Ala Ala Tyr Cys Leu
 1635 1640 1645
 Ser Thr Gly Cys Val Val Ile Val Gly Arg Ile Val Leu Ser Gly Lys
 1650 1655 1660
 Pro Ala Ile Ile Pro Asp Arg Glu Val Leu Tyr Gln Glu Phe Asp Glu
 1665 1670 1675 1680
 Met Glu Glu Cys Ser Gln His Leu Pro Tyr Ile Glu Gln Gly Met Met
 1685 1690 1695
 Leu Ala Glu Gln Phe Lys Gln Lys Ala Leu Gly Leu Leu Gln Thr Ala
 1700 1705 1710
 Ser Arg His Ala Glu Val Ile Thr Pro Ala Val Gln Thr Asn Trp Gln
 1715 1720 1725
 Lys Leu Glu Val Phe Trp Ala Lys His Met Trp Asn Phe Ile Ser Gly
 1730 1735 1740
 Ile Gln Tyr Leu Ala Gly Leu Ser Thr Leu Pro Gly Asn Pro Ala Ile
 1745 1750 1755 1760
 Ala Ser Leu Met Ala Phe Thr Ala Ala Val Thr Ser Pro Leu Thr Thr
 1765 1770 1775

Gly Gln Thr Leu Leu Phe Asn Ile Leu Gly Gly Trp Val Ala Ala Gln
 1780 1785 1790
 Leu Ala Ala Pro Gly Ala Ala Thr Ala Phe Val Gly Ala Gly Leu Ala
 1795 1800 1805
 Gly Ala Ala Ile Gly Ser Val Gly Leu Gly Lys Val Leu Val Asp Ile
 1810 1815 1820
 Leu Ala Gly Tyr Gly Ala Gly Val Ala Gly Ala Leu Val Ala Phe Lys
 1825 1830 1835 1840
 Ile Met Ser Gly Glu Val Pro Ser Thr Glu Asp Leu Val Asn Leu Leu
 1845 1850 1855
 Pro Ala Ile Leu Ser Pro Gly Ala Leu Val Val Gly Val Val Cys Ala
 1860 1865 1870
 Ala Ile Leu Arg Arg His Val Gly Pro Gly Glu Gly Ala Val Gln Trp
 1875 1880 1885
 Met Asn Arg Leu Ile Ala Phe Ala Ser Arg Gly Asn His Val Ser Pro
 1890 1895 1900
 Thr His Tyr Val Pro Glu Ser Asp Ala Ala Ala Arg Val Thr Ala Ile
 1905 1910 1915 1920
 Leu Ser Ser Leu Thr Val Thr Gln Leu Leu Arg Arg Leu His Gln Trp
 1925 1930 1935
 Ile Ser Ser Glu Cys Thr Thr Pro Cys Ser Gly Ser Trp Leu Arg Asp
 1940 1945 1950
 Ile Trp Asp Trp Ile Cys Glu Val Leu Ser Asp Phe Lys Thr Trp Leu
 1955 1960 1965
 Lys Ala Lys Leu Met Pro Gln Leu Pro Gly Ile Pro Phe Val Ser Cys
 1970 1975 1980
 Gln Arg Gly Tyr Arg Gly Val Trp Arg Gly Asp Gly Ile Met His Thr
 1985 1990 1995 2000
 Arg Cys His Cys Gly Ala Glu Ile Thr Gly His Val Lys Asn Gly Thr
 2005 2010 2015
 Met Arg Ile Val Gly Pro Arg Thr Cys Arg Asn Met Trp Ser Gly Thr
 2020 2025 2030

Phe Pro Ile Asn Ala Tyr Thr Thr Gly Pro Cys Thr Pro Leu Pro Ala
 2035 2040 2045
 Pro Asn Tyr Lys Phe Ala Leu Trp Arg Val Ser Ala Glu Glu Tyr Val
 2050 2055 2060
 Glu Ile Arg Arg Val Gly Asp Phe His Tyr Val Ser Gly Met Thr Thr
 2065 2070 2075 2080
 Asp Asn Leu Lys Cys Pro Cys Gln Ile Pro Ser Pro Glu Phe Phe Thr
 2085 2090 2095
 Glu Leu Asp Gly Val Arg Leu His Arg Phe Ala Pro Pro Cys Lys Pro
 2100 2105 2110
 Leu Leu Arg Glu Glu Val Ser Phe Arg Val Gly Leu His Glu Tyr Pro
 2115 2120 2125
 Val Gly Ser Gln Leu Pro Cys Glu Pro Glu Pro Asp Val Ala Val Leu
 2130 2135 2140
 Thr Ser Met Leu Thr Asp Pro Ser His Ile Thr Ala Glu Ala Ala Gly
 2145 2150 2155 2160
 Arg Arg Leu Ala Arg Gly Ser Pro Pro Ser Met Ala Ser Ser Ser Ala
 2165 2170 2175
 Ser Gln Leu Ser Ala Pro Ser Leu Lys Ala Thr Cys Thr Ala Asn His
 2180 2185 2190
 Asp Ser Pro Asp Ala Glu Leu Ile Glu Ala Asn Leu Leu Trp Arg Gln
 2195 2200 2205
 Glu Met Gly Gly Asn Ile Thr Arg Val Glu Ser Glu Asn Lys Val Val
 2210 2215 2220
 Ile Leu Asp Ser Phe Asp Pro Leu Val Ala Glu Glu Asp Glu Arg Glu
 2225 2230 2235 2240
 Val Ser Val Pro Ala Glu Ile Leu Arg Lys Ser Arg Arg Phe Ala Arg
 2245 2250 2255
 Ala Leu Pro Val Trp Ala Arg Pro Asp Tyr Asn Pro Pro Leu Val Glu
 2260 2265 2270
 Thr Trp Lys Lys Pro Asp Tyr Glu Pro Pro Val Val His Gly Cys Pro
 2275 2280 2285

Leu Pro Pro Pro Arg Ser Pro Pro Val Pro Pro Pro Arg Lys Lys Arg
 2290 2295 2300
 Thr Val Val Leu Thr Glu Ser Thr Leu Ser Thr Ala Leu Ala Glu Leu
 2305 2310 2315 2320
 Ala Thr Lys Ser Phe Gly Ser Ser Ser Thr Ser Gly Ile Thr Gly Asp
 2325 2330 2335
 Asn Thr Thr Thr Ser Ser Glu Pro Ala Pro Ser Gly Cys Pro Pro Asp
 2340 2345 2350
 Ser Asp Val Glu Ser Tyr Ser Ser Met Pro Pro Leu Glu Gly Glu Pro
 2355 2360 2365
 Gly Asp Pro Asp Leu Ser Asp Gly Ser Trp Ser Thr Val Ser Ser Gly
 2370 2375 2380
 Ala Asp Thr Glu Asp Val Val Cys Cys Ser Met Ser Tyr Ser Trp Thr
 2385 2390 2395 2400
 Gly Ala Leu Val Thr Pro Cys Ala Ala Glu Glu Gln Lys Leu Pro Ile
 2405 2410 2415
 Asn Ala Leu Ser Asn Ser Leu Leu Arg His His Asn Leu Val Tyr Ser
 2420 2425 2430
 Thr Thr Ser Arg Ser Ala Cys Gln Arg Gln Lys Lys Val Thr Phe Asp
 2435 2440 2445
 Arg Leu Gln Val Leu Asp Ser His Tyr Gln Asp Val Leu Lys Glu Val
 2450 2455 2460
 Lys Ala Ala Ala Ser Lys Val Lys Ala Asn Leu Leu Ser Val Glu Glu
 2465 2470 2475 2480
 Ala Cys Ser Leu Thr Pro Pro His Ser Ala Lys Ser Lys Phe Gly Tyr
 2485 2490 2495
 Gly Ala Lys Asp Val Arg Cys His Ala Arg Lys Ala Val Ala His Ile
 2500 2505 2510
 Asn Ser Val Trp Lys Asp Leu Leu Glu Asp Ser Val Thr Pro Ile Asp
 2515 2520 2525
 Thr Thr Ile Met Ala Lys Asn Glu Val Phe Cys Val Gln Pro Glu Lys
 2530 2535 2540

Gly Gly Arg Lys Pro Ala Arg Leu Ile Val Phe Pro Asp Leu Gly Val
 2545 2550 2555 2560
 Arg Val Cys Glu Lys Met Ala Leu Tyr Asp Val Val Ser Lys Leu Pro
 2565 2570 2575
 Leu Ala Val Met Gly Ser Ser Tyr Gly Phe Gln Tyr Ser Pro Gly Gln
 2580 2585 2590
 Arg Val Glu Phe Leu Val Gln Ala Trp Lys Ser Lys Lys Thr Pro Met
 2595 2600 2605
 Gly Phe Ser Tyr Asp Thr Arg Cys Phe Asp Ser Thr Val Thr Glu Ser
 2610 2615 2620
 Asp Ile Arg Thr Glu Glu Ala Ile Tyr Gln Cys Cys Asp Leu Asp Pro
 2625 2630 2635 2640
 Gln Ala Arg Val Ala Ile Lys Ser Leu Thr Glu Arg Leu Tyr Val Gly
 2645 2650 2655
 Gly Pro Leu Thr Asn Ser Arg Gly Glu Asn Cys Gly Tyr Arg Arg Cys
 2660 2665 2670
 Arg Ala Ser Gly Val Leu Thr Thr Ser Cys Gly Asn Thr Leu Thr Cys
 2675 2680 2685
 Tyr Ile Lys Ala Arg Ala Ala Cys Arg Ala Ala Gly Leu Gln Asp Cys
 2690 2695 2700
 Thr Met Leu Val Cys Gly Asp Asp Leu Val Val Ile Cys Glu Ser Ala
 2705 2710 2715 2720
 Gly Val Gln Glu Asp Ala Ala Ser Leu Arg Ala Phe Thr Glu Ala Met
 2725 2730 2735
 Thr Arg Tyr Ser Ala Pro Pro Gly Asp Pro Pro Gln Pro Glu Tyr Asp
 2740 2745 2750
 Leu Glu Leu Ile Thr Ser Cys Ser Ser Asn Val Ser Val Ala His Asp
 2755 2760 2765
 Gly Ala Gly Lys Arg Val Tyr Tyr Leu Thr Arg Asp Pro Thr Thr Pro
 2770 2775 2780
 Leu Ala Arg Ala Ala Trp Glu Thr Ala Arg His Thr Pro Val Asn Ser
 2785 2790 2795 2800

Trp Leu Gly Asn Ile Ile Met Phe Ala Pro Thr Leu Trp Ala Arg Met
 2805 2810 2815
 Ile Leu Met Thr His Phe Phe Ser Val Leu Ile Ala Arg Asp Gln Leu
 2820 2825 2830
 Glu Gln Ala Leu Asn Cys Glu Ile Tyr Gly Ala Cys Tyr Ser Ile Glu
 2835 2840 2845
 Pro Leu Asp Leu Pro Pro Ile Ile Gln Arg Leu His Gly Leu Ser Ala
 2850 2855 2860
 Phe Ser Leu His Ser Tyr Ser Pro Gly Glu Ile Asn Arg Val Ala Ala
 2865 2870 2875 2880
 Cys Leu Arg Lys Leu Gly Val Pro Pro Leu Arg Ala Trp Arg His Arg
 2885 2890 2895
 Ala Arg Ser Val Arg Ala Arg Leu Leu Ser Arg Gly Gly Arg Ala Ala
 2900 2905 2910
 Ile Cys Gly Lys Tyr Leu Phe Asn Trp Ala Val Arg Thr Lys Leu Lys
 2915 2920 2925
 Leu Thr Pro Ile Ala Ala Ala Gly Arg Leu Asp Leu Ser Gly Trp Phe
 2930 2935 2940
 Thr Ala Gly Tyr Ser Gly Gly Asp Ile Tyr His Ser Val Ser His Ala
 2945 2950 2955 2960
 Arg Pro Arg Trp Phe Trp Phe Cys Leu Leu Leu Leu Ala Ala Gly Val
 2965 2970 2975
 Gly Ile Tyr Leu Leu Pro Asn Arg
 2980

<210> 3
 <211> 38
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 Primer

<400> 3
 acgcgtctgc agcttaatgg cccaggacgc gatgcttg

<210> 4
<211> 36
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
Primer

<400> 4
acgcgtagat cttaccaagt gcgcaattcc tcgggg

36

<210> 5
<211> 34
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
Primer

<400> 5
tcagttggat agcgtcgacg ccggcaaata gcag

34

<210> 6
<211> 34
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
Primer

<400> 6
cgtcgacgcg atccaactga tcaacaccaa cggc

34

<210> 7
<211> 26
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic

Primer

<400> 7

cttgtaccat caattacacc atattc

26

<210> 8

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
Primer

<400> 8

gatagtgcc a atgcctatac ggg

23

<210> 9

<211> 41

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
Primer

<400> 9

cgtataggca ttggcactat ccttgtagca tcaattacac c

41

<210> 10

<211> 41

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
Primer

<400> 10

gggtgaattg atgggtacaag gatagtgcc a atgcctatac g

41

<210> 11

<211> 41

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
Primer

<400> 11

ttttttttgc ggccgcatgg tggggaactg ggcgaaggtc c

41

<210> 12

<211> 42

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
Primer

<400> 12

acgcgtaagc ttctattact cggacctgtc cctgtcttcc ag

42

INTERNATIONAL SEARCH REPORT

Intern: al Application No
PCT/US 00/25987

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/40 C12N5/10 C12N7/01 C07K14/18 C07K16/10
A61K31/70 A61K39/29

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, BIOSIS, EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	FORNS X. ET AL.: "Characterization of modified Hepatitis C virus E2 protein expressed on the cell surface." VIROLOGY, vol. 274, 15 August 2000 (2000-08-15), pages 75-85, XP002156408 the whole document	1-60
X	WO 96 40764 A (US HEALTH) 19 December 1996 (1996-12-19) abstract page 10, line 27 - line 32	1,10,12, 14,16, 18,19, 21,23, 25-27, 29,31, 32,34, 35,37-39
	--- -/-	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *8* document member of the same patent family

Date of the actual completion of the international search

4 January 2001

Date of mailing of the international search report

24/01/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Galli, I

INTERNATIONAL SEARCH REPORT

Intern: al Application No
PCT/US 00/25987

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 99 04008 A (US HEALTH) 28 January 1999 (1999-01-28) abstract sequence of E2 from isolate H77C ----	1-60
A	WO 94 26306 A (CHIRON CORP) 24 November 1994 (1994-11-24) abstract ----	1-60
A	MALET I ET AL: "YELLOW FEVER 5' NONCODING REGION AS A POTENTIAL ELEMENT TO IMPROVE HEPATITIS C VIRUS PRODUCTION THROUGH MODIFICATION OF TRANSLATIONAL CONTROL" BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, ACADEMIC PRESS INC. ORLANDO, FL, US, vol. 253, no. 2, 18 December 1998 (1998-12-18), pages 257-264, XP000961230 ISSN: 0006-291X the whole document ----	3-9
A	YI M. ET AL.: "Delineation of regions important for heteromeric association of Hepatitis C Virus E1 and E2." VIROLOGY, vol. 231, 1997, pages 119-129, XP002156409 the whole document -----	40-60

INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern. Application No

PCT/US 00/25987

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9640764 A	19-12-1996	US 6110465 A	29-08-2000
		AU 718503 B	13-04-2000
		AU 6157996 A	30-12-1996
		CA 2221313 A	19-12-1996
		EP 0832114 A	01-04-1998
WO 9904008 A	28-01-1999	US 6153421 A	28-11-2000
		AU 8488998 A	10-02-1999
		EP 0998567 A	10-05-2000
WO 9426306 A	24-11-1994	CA 2162557 A	24-11-1994
		EP 0697888 A	28-02-1996
		JP 8510240 T	29-10-1996

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER: _____**

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.